

or depressed line extends from the base of the serrature obliquely towards the base of the branch, and at its termination the surface of the branch is marked by a minute but distinct round tubercle.

This beautiful little species differs very distinctly from any others of this genus, in the thickened substance of its branches, the closely arranged serratures, and the minute tubercles at the base of the grooves or striæ. The specimens usually preserve considerable substance, and are far less flattened than most of the other species, owing either to their original character or to the nature of the surrounding matrix. The impressions of the oblique lines or striæ are often well preserved in imprints of the fossil left in the slate.

The impressions of *G. bryonoides* resemble those of this species; but the branches are broader, and the striæ are less rigid and less distinctly impressed, while the absence of tubercles, and the coarser serratures, when visible, at once serve to distinguish the species.

In mode of growth and general aspect this species resembles the *G. serratulus* (Pal. N. Y., vol. 1, p. 274, pl. 74, fig. 5, a, b.) of the Hudson River shales; but in the latter the serratures are coarser and more oblique, the lower side being much the longer. The branches of that species are also more distinctly linear, while in this they become gradually wider from the base, and are very distinctly striate and tuberculate in well-preserved specimens.

The preceding description applies to the specimens of this species where the branches diverge abruptly, or nearly at a right angle, from the radicle.

Locality and Formation.—Point Lévy, Hudson River Group.

Collector.—J. Richardson.

GRAPTOLITHUS BIFIDUS.

Description.—Two-branched; branches very gradually and uniformly diverging from the base to the extremities; surfaces obliquely striated; serratures moderately oblique; extremities often nearly vertical to the rachis, and submucronate (?); from thirty-eight to forty in the space of an inch; radicle short.

This species resembles in general features the *G. nitidus*, and might be mistaken for that species with the branches approximated by pressure. In several individuals examined the serratures are much closer, being from six to eight more in the space of an inch, while the general form is constant. The outer mar-