B

L

EUCEPHALUS MACOUNII, Greene, Pittonia, vol. IV, p. 70.

Along fences, Sea's Farm, near Victoria, Vancouver Island. Herb. No. 447. (John Macoun.) Distributed as Aster radulinus.

CENTAUREA SCABIOSA, L.

Along the Canadian Pacific Railway at Snellgrove, Ont. (Jas. White.) New to Canada and known from only one other locality in America. Determined by Dr. Robinson.

SENECIO OVINUS, Greene, Pittonia, vol. IV, p. 110.

S. resedifolius, Macoun, Cat. Can. Plants, vol. 1, p. 267 in part.

Mountain slopes, western summit of North Kootanie Pass, Rocky Mts., 1883. (Dr. G. M. Dawson.) High slopes of Sheep Mountain, Waterton Lake, Rocky Mts. Herb. No. 11,619. (John Macoun.) Described from the Sheep Mountain specimens.

VACCINIUM NIGRUM, Britt.

V. corymbosum, var. pallidum, Macoun, Cat. Can. Plants, vol. 1, p. 291.

Point Pleasant, N.S.; Englishtown, Cape Breton Island, N.S.; common in the vicinity of Ottawa, Ont., and at Niagara, Ont. (John Macoun.)

Lysimachia vulgaris, L.

Well established on Toronto Island, Ont. (W. Scott.)
Only Canadian record.

STEIRONEMA LANCEOLATUM, Gray; Macoun Cat. Can. Plants, vol. 1, p. 313.

Recorded from Ontario, but such specimens as we have seen so named are S. quadriflorum, Hitchc.

ACERATES LONGIFOLIA, Ell.

Dry sandy soil, southwest of Sandwich, Ont., 1893. (Alex. Wherry.) Our only Canadian specimens. The specimens referred here, Macoun, Cat. Can. Plants, vol. 1, p. 563, are A. viridiflora var. lanceolata, Gr.