

CAPPARIDACEÆ.

common in *STERIPHOMA ELLIPTICA*, *Spreng.* Hab. S. America.

RESEDACEÆ.

From the islanders about the *RESEDA ODORATA*, *Spreng.* "Mignonette." Cultivated. Hab. Egypt.

FLACOURTIACEÆ.

the northern as *FLACOURTIA PRUNIFOLIA*.
India, China, temperate regions

HYPERICACEÆ.

HYPERICUM — ? Pembroke Marshes.

VISMIA GUIANENSIS, *D. C.* Hab. Guiana.

CARYOPHYLLACEÆ.

DIANTHUS — ? "Pink." "Carnation." Varieties cultivated.

common weed *SAPONARIA CALABRICA*, *Guss.* "Soapwort." Hab. Calabria.

PORTULACEÆ.

PORTULACA OLERACEA, *L.* "Purslane" "Pursley." Hab.

common every size, almost south shore. It is used as All tropical as well as temperate regions. A common weed, generally found in cultivated ground. In flower end of March and beginning of April. It is sometimes used as a vegetable, boiled, and seasoned with pepper and salt. Pigs and poultry are fond of it.

FICOIDEÆ.

cultivated. *SESUVIUM PORTULACASTRUM*, *L.* "Sea Purslane." In sandy rocks below mud above high water mark. This plant is found on all temperate as well as tropical shores throughout the globe. Varieties

MESEMBRYANTHEMUM GLACIALE, *Haw.* "Ice Plant." Cultivated. Hab. Greece.

MALVACEÆ.

seed growing

MALVA — ?

MODIOLA CAROLINIANA, *G. Don.* (*M. multifida*, *Mænench*—*Malva caroliniana*, *L.*) Hab. N. America.