

***Acmaeops proteus* Ky.**

Yukon Territory; lat. $62^{\circ} 31' - 63^{\circ} 06' N.$; long. $137^{\circ} 30' - 139^{\circ} 30' W.$, 1913; D. D. Cairnes, collector, 6 specimens.

***Acmaeops pratensis* Laich.**

Yukon Territory, latitude and longitude as above; 1916, D. D. Cairnes, collector, 1 specimen. Near Sixtymile river, along meridian $141^{\circ} W.$, 1907, T. P. Reilly, collector, 1 specimen. Rampart House (18 miles south), Yukon Territory, D. H. Nelles, collector, 1 specimen. Fort Cudahy, Yukon Territory, 25, VIII, 1896, W. Ogilvie, collector, 2 specimens.

Genus ***Callidium* Fabr.*****Callidium subopacum*, n. sp.**

Length, 9 mm.: body piceous, above dull greenish, antennae reddish except first segment very slender; above subopaque; head rather obscurely punctured and reticulate, punctures coarser and more distinct behind; median line finely impressed, a wide transverse impression between the eyes; the pronotum much wider than long, nearly as wide as the elytra, widest about the middle, the sides broadly arcuately narrowed in front and more strongly narrowed behind the middle, the sides spongy from exceeding by dense, very deep, subcircular, margined punctures, with rather numerous long brownish hairs, a median amphora-shaped area impressed, the margins rather indefinite, nearly smooth, finely reticulate, the punctures coarse but very shallow, the pubescence red, short and erect, longer in front; the elytra thin, coarsely irregularly punctured, margins of the punctures indefinite, with minute setose punctures in their depth, these feebly granulate only towards the base.

Type, south of Rampart House, Yukon Territory, D. H. Nelles, collector; one paratype, same labels. Two other allied undescribed species are in our collection from southern British Columbia.

***Tetropium cinnamopterum* Ky.**

Fort Cudahy, Yukon Territory, 25-VIII-96, W. Ogilvie, collector. 4 specimens.

Length, 9-10 mm.; slender, piceous below, head and pronotum black, elytra smoky brown; pronotum subangularly rounded on sides, closely punctured and granulate on the sides, smooth, shining, and very sparsely punctured over nearly whole of disc, with a broad and strong anterior impression; the antennae rather slender.

FAMILY BUPRESTIDAE.

Only one specimen of this Family was included in the collection:

***Melanophila longipes* Say.**

Near Langton bay, Northwest Territories, 1910-11, R. M. Anderson, collector, 1632.

Other northern records in our collection: Near Sixtymile river, along 141^{st} meridian, Yukon Territory, 1907, T. P. Reilly, collector, 1 specimen; Stewart river, Yukon Territory, 1909, D. H. Nelles, collector, 1 specimen.

Occurs throughout Canadian forests from Nova Scotia to British Columbia and the Yukon.