

October 28th, patient was in good health and both sides of the neck were normal in appearance.

Tuberculosis of the Skin.

Miss J. M., aged 34 years, came to the Montreal Dispensary, May 23rd, 1905, and was referred to me by Dr. J. M. Jack, to whom I am indebted for having so patiently followed this case as well as several others.

Patient had an eruption on her nose and face, which began nine years ago. During that time she had a varied treatment, including the Finsen light, and X-rays for four months without benefit. The skin lesion was on the right temple, both cheeks and nose. There was some loss of tissue of the left nostril and some scarring on the right cheek. The skin was raised, indurated, and here and there a yellow spot appeared on which a crust had formed. She received her first injection of twenty-four grains on May 29th, 1905. By June 1st a change in the diseased area had taken place. The skin had become soft and pliable. There were numerous small yellow spots. These were tubercles which, upon microscopic examination, showed that they had undergone fatty degeneration in the centre. By June 5th the redness in the skin and induration were much less, and the yellow spots were beginning to dry up, and by June 11th were all absorbed. In some places the tissue began to whiten out. Before the second injection, June 12th, one or two tubercles underwent resolution, but after it a great many more were acted on. The area of diseased tissue steadily decreased. After the third injection only a few tubercles were acted on. She was injected July 31st, October 30th, and November 27th. Following each treatment the induration steadily diminished, and a few tubercles underwent resolution. As the tubercles disappeared the skin became paler and finally white scar tissue remained. At present time, February 16th, 1906, the disease is practically eradicated.

Case V.—Mr. C., came to the Montreal Dispensary and was referred to me for treatment by Dr. J. M. Jack. He complained of cough, loss of flesh and expectoration of purulent character, and numerous small ulcers on the right shoulder joint, which had been increasing in area during the last five years. There was no history of lues. Consolidation in right apex of lung. Skin on the right shoulder was raised, indurated, and here and there were numerous small ulcers, one as large as a fifty-cent piece. These ulcers had a crater-like appearance with overhanging edges and unhealthy base. Some of them