

TABULATED RESULTS—continued.

Case	Name	Sex	Age	Refraction.	Vision.	Deviation.	Degrees.	Abduction.	Adduction.	Symptoms.	Operation.	Result.
22	Miss B. ...	F.	20	R. — 10 D. L. — 9 D.		Hyper- exophoria	3 ex. 13	18°	34°	Frequent and intense headaches; new work always brings them on	Tenot. R. inf., L. ext. and L. sup. recti	Greatly relieved.
23	M. W. ...	F.	17	R. — 4.50 L. + 5.00		Hyper- phoria	3	16°	40°	In reading the print is unsteady, and this always causes headaches	Tenot. L. inf. and R. int. recti	Could use the eyes with perfect comfort a year later.
24	L. H. ...	F.	34	R. + 1.25 L. + 1.25		Hyper- exophoria	2 1/2 ex. 6	12°	17°	Asthenopia ...	Tenot. R. sup. and L. ext. recti	Much relieved.
25	I. C. ...	F.	19	R. — 2.25 L. — 2.25		Exophoria	6	18°	16°	Asthenopia and head- aches	Tenot. both ext. recti	Complete relief.
26	A. S. ...	M.	30	R. Emmetropia L. 55° + 1 D. ...		Exophoria	7	13°	15°	Blepharitis. Asthe- nopia	Tenot. both ext. recti	Much relieved.
27	W. L. W.	M.	35	R. 65° + 0.75 + 7.25 L. + 1.75		Hyper- phoria	12	6°	14°	Asthenopia ...	Tenot. R. int. rec- tus, later L. inf. rectus	Greatly relieved.
28	M. McG.	M.	18	R. + 1.50 L. — 1.50		Esophoria	6	5°	65°	Asthenopia and chronic conjunctiv- itis	Tenot. both int. recti	Complete relief.
29	A. H. ...	M.	20	R. 30° — 1.75 — 0.75 L. 90° — 1.50 — 1.00...		Esophoria	8	6°	65°	Asthenopia conjunc- tivitis. Hyper- emia not relieved by glasses	Tenot. int. recti at interval of 2 weeks	Complete relief.
30	Mrs. F. ...	F.	42	R. + 1.00 L. + 1.00		Exophoria	10	12°	17°	Persistent headaches for many years	Tenot. both ext. recti at interval of 5 days	Eight months later reported much better.

31	H. L. ...	M.	16	R. + 1.25 L. + 1.25		Esophoria	11	5°	63°	Headaches induced by any new work	Tenot. both int. recti at intervals	Reported that the eyes could be used with
----	-----------	----	----	------------------------	------------	-----------	----	----	-----	--------------------------------------	--	--