

Cranioscopy. Fig. 90. No. 9.

Age, 65
Sutures, serrations very complicated
Pterion, H, but on left side bridge very narrow
Artificially lengthened occipital foramen comes
to E
Lateral parietes, bulging
Glabella, very prominent.
Superciliary ridges, very prominent.
Frontal eminences, small.
Forehead, retreating
Median line, presents at 2 centimetres above
lambde a rounded arrivers above
lambda a rounded eminence about 2 cen-
timetres in diameter
Probola, not prominent.
Sub-iniac curve, level with condyles
Mastoid process, very long, thick and rough,
one is 30.8 millimetres in length
Malar bones, very thick, with muscular attach-
ments strongly marked
Hollow at root of nose, shallow.
Inferior border nares, heart shaped.
Inferior border orbit, much thickened
Muscular attachments, very strongly marked on
all parts of the skull. Posterior root of
zygoma is raised behind into a prominent
ridge with sharply marked hollow behind
it. Similar markings are present on each
side in lambdoid suture about 2 centim.
above asterion, groove looking backward
and inward. On superior curved line is
another sharp depression looking unward
with a ridge below it

Cr	aniometry.
1	Millimetres
1	184
2	149
3	140
4	100
5	108
6	114
7	16
8	112
9	135
10	55
11	62
12	300
13	240
14	286
15	52 6
16	83
17	147
18	54
19	27
20	80.9
21	67.1
22	92.6
23	76.0