

b³. Ovipositor decidedly curved.

c¹. Size small, less than 14 mm.; tegmina without the two black lines usually present on either side of the speculum.....*Gracile*, Harr.

c². Size medium or large, at least 16 mm. long.

d¹. Face pale or unicolorous, never with reddish-brown or fuscous stripe down the middle.

e¹. Hind femora short, less than five-sixths the length of the body and not exceeding 15 mm. in length.

f¹. Tegmina not much exceeding the hind femora ; ovipositor less than 9 mm. long ; posterior margin of the lateral lobes strongly sinuate ; anterior and middle tibiae green or yellowish.....*Vulgare*, Harr.

f². Tegmina far surpassing the tips of the hind femora ; ovipositor more than 10 mm. long ; posterior margin of the lateral lobes of the pronotum distinctly but not strongly sinuate ; anterior and middle tibiae fuscous black.....*Robustum*, Red.

e². Hind femora longer, at least five-sixths as long as the body and not less than 17 mm. in length ; ovipositor less than half as long as the hind femora.

f¹. Pronotum longer, more than one-fourth as long as the body ; tegmina not far surpassing the tip of the hind femora.

g¹. Tegmina not narrowed in the middle ; anal cerci of the male slender, the basal tooth short and weak ; tegmina and wings olivaceous brown ; size less than medium.....*Campestre*, Blatchley.

g². Tegmina narrowed in the middle ; anal cerci of the male swollen, armed within with a strong basal tooth ; posterior margin of the lateral lobes of the pronotum strongly sinuate ; size greater than medium.....*Glaberrimum*, Burm.