

Muhl. in herb. an original one accidum. It is orbicular, plane plicate-papulose cunose; apoth. thin, coarctate. 321. *Muhl.* ! Ohio

3. L. *Burgessii*, Fr. Th. membranaceous, subimbricate, glaucous-fuscous, somewhat spongy and downy beneath, the lobes rounded, sinuate-laciniate, crisped and minutely lacerate-dentate at the margins; apoth. depressed; disk somewhat concave, fuscous, with an elevated, lacerate-dentate or foliose margin. *Collema*, Ach. Syn. p. 320. *Hook.* / Br. Fl. 2, p. 211. Icon, E. Bot. t. 300.

Trunks. Mountains of New England.

Tribe II. EPHEBIDEÆ.

III. EPHEBE, Fr.

Apothecia formed from the thallus, from concave becoming plane, and at length convex, black, the margin evanescent. Thallus filamentous, not gelatinous.

E. *pubescens*, Fr. Thallus decumbent, softish, terete, black, the branches entangled, capillaceous; apothecia of the same color. Fr. Fl. Scan. p. 294. *Cornicularia*, Ach.— β . *fibrillosa*, Ach.; thallus obscurely fuscous, smoothish, very delicate, branched, somewhat hirsute with numerous, flexuous, branched, subclavate fibres. *Cornicularia pubescens*, var. *fibrillosa*, Ach. Syn. p. 302.

Rocks and stones; — α , in alpine districts. Greenland, Dill. White Mountains. — β , North America, Ach.

Fr. Fl. Scan.

ey. Pennsyl-

, very thin and small, subimbricate (small) scattered margin. Fr. ! F. 3.

England. Ne