nctional eye disturb in the day, frequent sain in the arms, face and nose, a greenish On the 21st of May, ater being used; disonce appeared; her y flushings until the y clear; color coming fect; and her genetter.

very yellow cheeks, which extend down palpitation indicates on filled. Is losing er husband's poverty sicipations, and her ane. A daily flushgust, 1887, and conat they were taken February, 1888, she tely revolutionized and strong; they en; their presence better after taking a be upon her, she soundly thereafter. by great tympanve. Formerly had she never has it. he, with vomiting ours every two or not had one. Uriwhich lightens her

l chronic digestive years; has gastric dyspepsia characterized by pain, load, weight, gas acid and bitter pyrosis; could cat but few things, butter especially distressing her; constipated and costive since girlhood; for years had a sense of pressure in the rectum; at times, of late, has had it all the time. Flushing first given the 16th of May, 1887, twice per week, at bedtime. At first they produced pain and nausea, and once vomiting followed. These adverse symptoms were relieved when the water passed. The discharges from her bowels were simply enormous. They were continued until she passed yellow feces. On the 28th of June she was eating everything. She has become a good breakfaster. On the first of August she was doing all her own housework, the first time in three years.

Case IV.—Miss C., age 30, always constipated. muddy complexion, chloasmic spots all over her face, capricious appetite, seeks relief for a progressively increasing nervousness, which has been upon her for the past three years, and is very much worse in the past six months, and threatens to lead her to a suicide's grave. She experiences sudden misgivings; she has nervous prostration, colon filled to an enormous extent. Flushing was used and continued daily for a period of one month, when she reported that she was better in every way, does not give out any more; is stronger in all things; her complexion is clearing out; the chloasmic spots are unchanged; the appetite decidedly vigorous. In this case the fecal reliefs were enormous, and the improvements so rapid that there was no question of there being cause and effect.

Case V.—Mrs. A., age 34, has been having epileptic convulsions since April, 1883, two years subsequent to her only confinement. Immediately after her confinement, in February, 1881, she began to experience nervous symptoms, such as numbness, twitching of the left side of the body; these increased until convulsions began in April, 1883. From that time until the 1st of May, 1887,