

CAMPANULACEÆ.

Campanula rotundifolia

N.B. — The Newfoundland variety (?) of this plant differs from the European form in being much less branched, with fewer, but much larger flowers; average specimens of which measure 1 inch long by 1 inch 2 lines wide, and are much more pentagonal in form. The plant is called 'thimble-flower' in Newfoundland.

ERICACEÆ.

Gaylussacia resinosa

Vaccinium cespitosum

" Canadense

" Pennsylvanicum

" corymbosum

Oxycoccus palustris

" macrocarpus

Chiogenes hispidula

Kalmia glauca

Epigaea repens

Arctostaphylos Uva-ursi

? Gaultheria procumbens

Andromeda polifolia - glaucophylla

" calyculata

Ledum palustre - ?

Pyrula secunda

" minor

" chlorantha

Moneses grandiflora

? Chimaphila umbellata ?

Monotropa uniflora

" Hypopitys

Calluna vulgaris

N.B. — Through the kindness of the Bishop of N.F.L., I have recently received a liberal supply of this plant in flower. The specimens were gathered near Capelin Bay in about 47° N.L., and 53° West long. All the known localities for this plant in N.F.L. appear to be in about the same latitude, and between 53°—54° west longitude! viz., "Head of St Mary's Bay and Trepassoy Bay," (Mr. W. E. Cormack).

Professor Geo. Lawson, in Transactions of Nova Scotia Institute of Nat. Science records having gathered Calluna vulgaris on Ulston Farm, St. Anns Bay, Cape Breton Island. It has also been gathered at Tewksbury, Massachusetts.

PRIMULACEÆ.

Primula Mistassinica	} both common.
" farinosa	
Trientalis Americana	
Samolus valerandi ?	

PLANTAGINACEÆ.

Plantago major (Introduced?)
" lanceolata - decurva

PLUMBAGINACEÆ.

Statice Limonium
Arméria vulgaris (rare?)

LENTIBULACEÆ.

Pinguicula vulgaris	
Utricularia intermedia	
* " minor	} rare?
" vulgaris	
" ? subulata	
" cornuta	

OROBANCHACEÆ.

Conopholis Americana
Aphyllon uniflora

SCROPHULARIACEÆ.

Veronica Anagallis: americana
" scutellata
" serpyllifolia
Castilleja septentrionalis
Pedicularis Canadensis ?
Rhinanthus Crista-galli
Euphrasia officinalis, common form.
" purpurea, n. sp.? This form of eyebright occurs in considerable patches on the sea coast at Cow Head, N.F.L., and is readily distinguished when growing by its being much smaller proportionally in all its parts, and by its dark purple flowers. (E. gracilis? Fries.)