

**Pneumonia, Broncho-cath-
rhal, or lobular** *(continued)*

- Fever in, I. 669, 673
- Hygienic treatment in, I. 677
- Hypophosphite of lime in, I. 677
- Indications for treatment of, I. 667-670
- Inhalation of oxygen in, I. 676
- Iodine in, I. 677
- Ipecacuanha in, I. 671
- Leeches in, I. 675
- Local applications of cold in, I. 674-75
- Opium in, I. 673, 676
- Pyrexia of, I. 674
- Quinine in, I. 674
- Sprays in, I. 671
- Stimulating embrocations in, I. 673
- Strychnine in, I. 676
- Symptoms and physical signs in, I. 669-70
- Tendencies in severe cases, I. 669
- Tincture of aconite in, I. 673
- Tonica in, I. 677
- Turpentine inhalation in, I. 673
- Vomiting in, I. 675
- Warm saline drinks in, I. 672
- See also Additional Formulae, I. 679-80*

Pneumonias, Secondary, I. 678**Pneumothorax and Pyo-
pneumothorax, II. 68-70**

- Blood-letting in, II. 68
- Capillary trocar in, II. 69
- Injection of morphine in, II. 69, 70
- Operations in, II. 68-70
- Opium in, II. 68, 70
- Quinine in, II. 68, 70
- Stimulants in, II. 68

**Polio-myelitis, Acute. See Para-
lysis, Infantile, II. 292-98****Posterior sclerosis. See Ataxy.****Locomotor, II. 271-282****Pseudo-leukemia. See Hodg-
kin's disease, I. 515-18****Pseudo-membranous stoma-
titis. See Stomatitis, Ul-
cerative, I. 8-10****Pulmonary emphysema. See
Emphysema, I. 623-32****Pulmonary tuberculosis. See
Phthisis, I. 737-II. 77****Pyuria and Pyelitis, II. 197-200**

- Alkalies and salines in, II. 199
- Bacilli in, II. 198
- Balsamic stimulants in, II. 199
- Baths in, II. 199
- Causation of, II. 197
- In gouty and rheumatic subjects, II. 199
- Mineral waters in, II. 199 ;
- Quinine in, II. 200

Pyuria and Pyelitis *(continued)*

- Surgical treatment in, II. 200
- Symptoms of, II. 198
- Turpentine in, II. 199
- Urinary antiseptics in, II. 199
- See also Additional Formulae, II. 204*

**Quinsy. See Tonsillitis, Acute,
I. 14-20****Rachitis, or Rickets, II. 542-48**

- Arsenic in, II. 548
- Carbonate of lime with salicylate of bismuth in, II. 548
- Causes of, II. 543
- Cod-liver oil in, II. 546, 548
- Dietetic measures in, II. 540-47
- Hygienic measures in, II. 547
- Indications for treatment of, II. 545-46
- Iron in, II. 548
- Phosphorus in, II. 548
- Prevention of deformity in, II. 547-48
- Sodium bromide with chloral in, II. 548
- Symptoms of, II. 543-45
- See also Additional Formulae, II. 556*

**Rectal feeding, General remarks on,
I. 41-47**

- Nutrient enemata and suppo-
sitories in, I. 42-46
- Peptonized food in, I. 41-42

**Remittent malarial fever. See
Malarial fevers, II. 726-44****Renal calculi, II. 187-90**

- Causation of, II. 189
- Composition of, II. 188
- Symptoms and consequences of, II. 189
- See also Hydronephrosis, II. 195-97 ; Pyuria and pyelitis, II. 197-200, and Renal colic II. 190-94*

Renal colic, II. 190-94

- Alcohol in, II. 191
- Alkaline drinks in, II. 193
- Causation of, II. 190-91
- Chloroform inhalations in, II. 192
- Henbane in, II. 192
- In children, II. 194
- Indications for treatment of, II. 191
- Injections of morphine and atro-
pine in, II. 191
- Mineral waters in, II. 193
- Operative measures in, II. 194
- Opium in, II. 192
- Paroxysm in, II. 190
- Piperazin in, II. 194