

Athasanus striola Fall. Occurred abundantly near South Falls, on a low meadow overgrown with swamp grasses ; here they were of large size, measuring in some cases 6 mm. to tip of the elytra ; elsewhere, scattering and of normal-size.

Allygus irroratus Say. Taken in moderate numbers. Both the normal form and the large variety, with the vertex produced and flattened, occurred together here as elsewhere ; the variety predominating in moist situations.

Jassus immistus Say. Common among low bushes and briars. This pretty insect seems to be congeneric with a form occurring near Buffalo, of which I have seen specimens in the National Museum labelled *Scophoideus jucundus* Uhl. A third and undescribed species occurred to me at Muskoka.

Platymetopius acutus Say. Moderately abundant.

Deltocephalus debilis Uhl. Rare. Occurs also at Buffalo.

Deltocephalus configuratus Uhl. Abundant here as elsewhere.

Deltocephalus sayi. Fitch. Scarce.

Deltocephalus inimicus Say. Very common on cultivated lands.

Typhlocybidæ.

Several species of these minute insects occurred ; but as the North American species have been but little studied they are omitted from the present list.

Psyllidæ.

Psylla carpini Fitch. Abundant on the blue beech.

Aphididæ.

Want of time precluded any attempt at collecting the *Aphididæ* and *Coccidæ*.

In addition to the 141 species here enumerated a number of doubtful forms were taken, belonging mostly to the Homoptera ; giving a total of something over 150 species as the fruits of ten days collecting.