

Section 4. Toes, 3; bill, under .60.*Aegialitis semipalmata.*

Bill, orange at base, the tip, black; legs, dull flesh color; a black stripe from bill passing under eye.

Semipalmated Plover.*Aegialitis semipalmata.*

See page 75.

Bill, orange at base, the tip, black; the legs, orange yellow; two middle tail feathers, tipped with white; no black stripe from bill to eye; black breast band not confluent. **Species not found west of the Rocky Mountains.**

Piping Plover.*Aegialitis meloda.*

See page 75.

Bill, orange at base, the tip, black; legs, orange yellow; middle tail feathers, tipped with white; no black stripe from bill to eye; **a continuous black band on breast.** **Species not found west of Rocky Mountains.**

Belted Piping Plover.*Aegialitis meloda circumcincta.*

See page 76.

*Aegialitis nivosus.*

Bill, entirely black; legs, slate color; two outer tail feathers, entirely white; two middle feathers, **not** tipped with white; no black stripe from bill to eye. Ranges from Texas and Kansas west to the Pacific Ocean; casual in Western Florida and Cuba; **not known to occur on the Atlantic Coast.**

Snowy Plover.*Aegialitis nivosus.*

See page 76.

Section 5. Toes, 3; bill, over .60.

A very small web between outer and middle toes; bill, large and thick; a band of black (male) or brown (female) on breast.

*Aegialitis Wilsonia.***Wilson's Plover.***Aegialitis wilsonia.*

See page 77.