

3. *L. insueta*, Guén.—Abundant on treacle and sumach blossoms, in July.

Distribution.—Florida and New York, (E. Doubleday); Nova Scotia, (Redman.)

4. *L. straminea*, Weit.—On treacle, beginning of July.

Distribution.—New York, (E. Doubleday), (Eu.)

Genus 2. NONAGRIA, Ochs.

1. *N?* *intractabilis*, Walker, *M.S.S.*, N. Sp. "*Fœm.* Albida cinerea, gracilis; palpi porrecti, extus nigricantes; caput sat superantes, articulo 3o. longo lanceolato; pedes breves, validi, sub testacei; alæ latiusculæ, non longæ, fusco nigroque conspersæ, lunulis marginalibus fuscis; anticæ testaceo subvariæ, lineis duabus albidis undulatis, valde indistinctis, orbiculari et reniformi albedo marginatis."

"*Female*. Whitish cinereous, slender. Palpi porrect, extending rather far beyond the head, blackish exteriorly; 3rd joint lanceolate, full half the length of the 2nd. Abdomen testaceous at the tip. Legs slightly testaceous, short, robust; spurs short stout. Wings rather broad, not long, minutely brown-and-black speckled; a marginal line of brown lunules. Forewings partly and indistinctly testaceous, somewhat rounded at the tips, with two very indistinct undulating whitish lines; orbicular and reniform marks whitish bordered, the former oblong the latter transverse and rather narrow. Length of the body $3\frac{1}{2}$ lines; of the wings 9 lines." Walker *M.S.S.*

Family, GLOTTULIDÆ, Guén.

Genus 1. EUDRYAS, Boisd.

1. *E. grata*, Fab., Notodonta? *grata* Harris. (Animals and Plants of Massa).—Not common.

Distribution.—New York (Emmons); Massachusetts (Harris); Sorel.

Family, APAMIDÆ, Guén.

Sub-Family, GORTYNIDES, Duponch.

Genus 1. Hydræcia, Guén.

1. *H. nictitans*, Linn.—Common end of July and beginning of August.

Distribution.—New York, (Doubleday); Mass., (Sheppard); Albany River, Hudson's Bay, (G. Barnston); Nova Scotia, (Redman); (Eu.)

2. *H. lorea*, Guén.—Abundant especially on blossoms of *Asclepias cornuti* in July.

Distribution.—New York, (E. Doubleday.)