

PRELIMINARY CATALOGUE OF THE ARCTIIDÆ OF TEMPERATE NORTH AMERICA, WITH NOTES.

BY JOHN B. SMITH, NEW BRUNSWICK, N. J.

(Continued from page 208, Volume xxii.)

H. roseata Wlk.1866—Wlk., App. to Lord's Trav. in Vanc., 336 (?), *Halisidota cinnamomea* Bdv.1868—Bdv., Lep. Cal. (Ann. Soc., Belg., XII.), 80, *Phægoptera*.1869—Grt. & Rob., Trans. Ann. Ent. Soc., III., 175, *pr. syn.*

Habitat—Vancouver, Calif.

As the description of this species is not readily accessible to students, I reproduce it here :—

"Female rosy red; body densely clothed, and partly pale yellow beneath; head with a pale yellow band on the front; palpi extremely short; thorax with six longitudinal pale yellow streaks; abdomen rosy, lanuginous and partly yellow to the base, extending much beyond the hind wings; fore wings with some pale yellowish streaks toward the base, with three exterior whitish macular, very oblique bands; spots mostly cuneiform; costa straight; tips slightly acute; exterior borders slightly convex, extremely oblique; first and second inferior veins contiguous at the base; third very near the second; fourth remote from the third; hind wings whitish cinereous, slightly hyaline; veins and fringe slightly yellowish. Length of the body, 7 lines; of the wings, 20 lines.

"This species may form a new genus. It differs somewhat from *Halisidota* in the structure of the veins of the fore wings."

H. scapularis Stretch.1885—Stretch, Ent. Amer., I., 106, *Halisidota*.

Habitat—New Mexico.

H. significans Hy. Edw.1888—Edw., Ent. Amer., III., 182, *Halisidota*.

Habitat—New Mexico.

H. sobrina Stretch.1873—Stretch, Zyg. and Bomb., 87, 135, pl. VI., f. 10, ♂, *Halisidota*.1873—Edw.*, Proc. Cal. Ac. Sci., V., 369, *Halisidota*.

Habitat—California.

Food-plant—*Pinus insignis*.