

each vine with an iron bar to the depth of a foot or more. Soot is also recommended, to be strewed around the vines.

It is stated that the insect is less injurious to vines grown on sandy soil ; also to those grown on lands impregnated with salt.

Since large numbers of these insects, both winged and wingless, are known to crawl over the surface of the ground in August and September, it has been suggested to sprinkle the ground about the vines at this period with quicklime, ashes, sulphur, salt or other substances destructive to insect life. The application of fertilizers rich in potash and ammonia have been found useful, such as ashes mixed with stable manure or sal-ammoniac.

A simple remedy for the gall-inhabiting type is to pluck the leaves as soon as they show signs of the galls, and destroy them.

Several species of predaceous insects prey on this louse.

A black species of Thrips with white fringed wings deposits its eggs within the gall, which, when hatched, produce larvæ of a blood red color, which play sad havoc among the lice. The larva of a Syrphus fly, *Pipiza radicum*, which feeds on the root louse of the apple, see figure 18, has also been found attacking the Phylloxera. Another useful friend is a small mite, *Tyroglyphus phylloxera*, P. & R., see fig. 19, which attacks and destroys the lice, and associated with this is sometimes found another species, *Hoplophora arctata*

Riley, of a very curious form, reminding one of a mussel. The lice are also preyed on by the larva of a Scymnus, a small dull colored lady bird ; also by several other species of the lady-bird family, and by the larvæ of lace-wing flies.




Fig. 18.




Fig. 19.