

Hab.—Morgantown, W. Va.

Bred May 4 and June 24, 1892, from *Anthaxia viridicornis* infesting Willow; and May 26, from an *Agrilus* larva living under the bark of Dogwood, *Cornus florida*.

(12) *C. pityophthori*, sp. n. *Caenophanes pityophthori*, Ashm. M. S., Ins. Life, IV., p. 258.

♂.—Length, 2 mm. Black; second abdominal segment at base and sometimes the petiole at apex ferruginous; head shining, faintly transversely aciculated on vertex; thorax opaque, minutely granulated; metathorax areolated, the pleura rugose; antennæ 22-jointed, black, the 3 basal joints yellow; abdomen elongate ovate, black, polished, the first two segments and the third at extreme base or the suture striated. Legs pale yellowish. Wings hyaline, the first abscissa of radius less than two-thirds the length of the second.

Hab.—Morgantown, W. Va.

Bred Feb. 24, 1892, from *Pityophthorus* sp. living under the bark of a small dying Spruce, *Abies nigra*.

(13) *C. hylotrupidis*, sp. n. *Caenophanes hylotrupidis*, Ashm., MS., Ins. Life, IV., p. 258.

♂, ♀.—Length, 2.5 to 3 mm.; ovipositor two-thirds the length of abdomen. Pale ferruginous; metathoracic sutures fuscous. Head above transversely aciculated; thorax finely granulated, the middle lobe posteriorly strongly rugose; metathorax, except the two basal areas, rugose, and with a Λ-shaped carina at the middle; first abdominal segment and basal two-thirds of the second striate, rest of abdomen smooth, polished, the second segment with a transverse impressed line at the middle. Antennæ in both sexes 25-jointed. Wings hyaline, the stigma and nervures pale brown, the first and second branches of the radius about equal in length.

Hab.—Morgantown, W. Va.

Bred April 4, 1891, from *Hylotrupes ligneus* living in *Juniperus virginiana*.

Subfamily HELCONINÆ.

HELCON, Nees.

(14) *H. occidentalis*, Cr. Helcon tetrapodii, Ashm., MS. (olim), Ins. Life, IV., p. 259.

Bred July 14, 1891, from *Tetrapodium cinnamoplerum* living in sapwood of Spruce log *Abies nigra*.