Caloptenus	(?) bilituratus, Walk.	Acridium alutaceum, Harr.
"	lurida, G. M. Dodge.	" emarginatum.
"	minor, Scudd.	Tropidacris dux, Scudd.
"	punctulatus, Uhler.	Tettix ornata, Scudd.
"	occidentalis, Thos.	" granulata, Scudd.
. "	regalis, Dodge, n. sp.	" cucullata, Scudd.
"	volucri, (?).	" triangularis, Scudd.
"	angustipennis, (?).	Tettigidea lateralis, Scudd.
Acridium a	mericanum, Scudd.	· " polymorpha, Scudd.
" a	mbiguum, Thos.	Batrachidea cristata, Scudd.

And in addition I have

3 species of Pezotettix, not yet determined.
2 " Caloptenus " "
2 " Œdipoda " "

And 3 others of different genera not yet determined, making a total of 95 species for Nebraska thus far.

TINEINA.

BY V. T. CHAMBERS, COVINGTON, KY.

BATRACHEDRA.

B. striolata? Zell. (Asychna? pulvella Cham.)

In "The Bulletin of the Geo. Survey," v. 3, p. 134, I have referred to two species or varieties of Batrachedra found by me in Colorado. Some of these specimens found at Colorado Springs I referred doubtfully to the European species B. præangusta. Others found higher up the mountains I named B. Clemensella, stating, however, the doubt whether the two forms were really distinct. After the greater portion of that paper was in the hands of the publisher, I received from Mr. Stainton two specimens of præangusta, by which I was enabled to recognise my B. Clemensella as a variety simply of that species, and to increase my doubts whether the supposed species were really distinct; for while Clemensella