

Therapeutic Notes.

FLATULENCE AND CHOLIC IN INFANTS.

R_y Magnesii carb..... gr. iiss
 Rhei pulv..... gr. ¼
 Syr. zingiberis..... m v.
 Aq. menth. pip..... q.s. ad 3 j

M. Sig. : Given every two to four hours to an infant three or four months old.—*Med. Rec.*

FATTY HEART.

A formula credited to A. Robin, by *Merck's Archives* is :

R_y Sodium arsenate..... ¼ gr.
 Potassium iodide ¾ gr.
 Powd. nux vomica..... ⅛ gr.
 Powd. rhubarb..... ¾ gr.
 Ext. dulcamara..... 1 ½ grs.

Make into one pill. Take one such daily.

Professor Pel, of Amsterdam, after a great deal of experience, has found that the following powder is a most excellent remedy for gastric acidity.

R_y Bicarbonate of soda 10.00 3 iiss
 Calcined magnesia 8.00 3 ii
 Bromid of soda..... 10.00 3 iiss
 Carbonate of bismuth..... 5.00 gr. lxxv
 Sugar of milk..... 10.00 3 ijss
 Oil of fennel..... .26 gt. iv

Mix. Dose, from half to a whole teaspoonful, to be taken an hour after eating. An extra dose in case of pain.
 —*Buffalo Med. Jour.*

AN INTESTINAL ANTISEPTIC.

A very good intestinal antiseptic is this which is credited by the *Med. News* to the *Jour. de Med. de Bordeaux* :

R_y Betanaphthol... 3.00 gr. xlv
 Chloroform03 gtt. xv
 Castor oil..... 90.00 3 iii
 Spirit of peppermint..... .02 gtt. v

Mix. Adult dose : A teaspoonful in port wine, beer or sweetened black coffee. Child's dose : A teaspoonful.