

Fig. 114. No. 47.

| Cranioscopy.  | Craniometry   |  |
|---|---|--|
| Both zygomatic proesses absent.  Age, 65  Sex, male.  Sutures, indistinct.  Pterion, H.  Inion, 2, rough.  Artificially lengthened occipital foramen, A.  Lateral parietes, protuberant.  Glabella, very prominent.  Superciliary ridges, very prominent.  Frontal eminences, indistinct.  Forehead, markedly retreating.  Median line, elevated from glabella to metopic point where it divides and encloses a rounded elevated triangular space which extends 3 centimetres on each side of bregma, sagittal suture raised.  Probola, medium.  Sub-iniac curve, does not extend below condyles Mastoid process, long, rough.  Hollow at root of nose, deep, nose has been | 1 2 3 4 5 6 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 | Millimetres. 190.5 138 144 94 104 105 10 112 119 77 44 295 236* 279 515 78 141 52 26 72.4 68.0 |
| broken in life, and is somewhat deflected.<br>Inferior border of nares, sides ascend rapidly  | 22 *<br>23  | 90.4<br>75.5   |

\*Over superciliary ridges=250 mm.

See page 60 for Key to Measurements.