THE CANADA MEDICAL RECORD.

Vot. XV.

MONTREAL, DECEMBER, 1886.

No. 3.

ORIGINAL COMMUNICATIONS.	
Rock and Rye in Acute and Subacute	
Bronchitis	-1
BB	49
SOCIETY PROCEEDINGS.	
Medico-Chirurgical Society of Mon-	
treal	5
PROGRESS OF SCIENCE,	
A Simple Method for the Diagnosis of Organic Valvular Diseases of the	
Organic Valvular Diseases of the	
Heart	5
Irritable Brain in Children	5
The Value of External Applications in	

		~
	the Treatment of ChildrenEffects of Cold Water	59
0	Bismuth Subnitrate in Burns	6:
9 9	Weaning. Insomnia in the Aged	6 6
	Winter Indigestion Biliousness	6
2	The Dietetics of Pulmonary Phthisis The Dietary in Indigestion,	6
	Sore Nipples.	6
	Camphor, Chloral and Cocaine in Footh- ache	6
5	The use and abuse of Tea Phylolacca Decandra in Treatment of	6
	Bronchocele	7

ĺ	CONTENT	S	5. · · ·	
And and an an an	the Treatment of Children.—Effects of Cold Water Bismuth Subnitrate in Burns	59 63 64 64 65 66 8 68 68 68 68 70	The Treatment of Bronchitis EDITORIAL. College of Physicians and Surgeons, Pro-	71 71 72 72 72 72 72 72
1	LIUNCHUUUIC	10	1	

Original Communications.

ROCK AND RYE IN ACUTE AND SUB-ACUTE BRONCHITIS.

BY HEBER BISHOP, B.A., C.M., M.D., BOSTON, MASS.

I have had excellent results in the administration of rock and rye in acute and sub-acute forms of bronchitis, and have found it of particular value in children. The ease with which the ordinary patient will take the drug renders it worthy of a trial.

As a proof of its efficacy I will cite one instance (that of a child two years old). I was called to the patient at 5 o'clock in the evening, found that the child had commenced with an ordinary coriza, running from the nose, which had been observed for two or three days ; the night before had commenced to cough, which during the day had become so severe that I was summoned in the afternoon.

The general appearance of the child was indicative of capillary bronchitis respirations 72 per minute, pulse 150, temperature 102° the alæ nasi dilated and the face turgid and purple with that frightfully distressed appearance. Urine scanty (had not micturated for 18 hours) ; with each cough the child would cry out.

Upon listening with the stethoscope coarse, harsh, rales were heard over both lungs, with some fine mucous crepitation over a portion of the left lung behind. I immediately ordered linseed meal poultices to be applied, enveloping the chest and back completely, and prescribed a mixture of syr. ipecac. (min. 5) syr. Acacia (min. 15) and Morphia (gr. r-50th.) to be administered every four or five

hours; and directed that a teaspoonful of rock & rye and glycerine be given in as much water every two hours. At my visit next morning found the child better and sleeping quietly, did not disturb Discontinued the syr. ipecac. Acacia and her. morphia but continued the rock & rye and glycerine at regular intervals of two hours.

At my evening visit found the child sitting up and playful, all acute symptoms subsided, pulse 118, temperature 99. "Respirations 33. This remarkable change had occurred in 24 hours.

In nearly every case of bronchitis that I have given it, rock & rye controlled the cough besides acting as a stimulant, and in young children it does not bind up the bowels the same as Brandy will often do. In winter cough (so called) it exerts a very soothing effect.

UNIVERSITY BISHOP'S COLLEGE.

THE ANNUAL DINNER OF THE MEDICAL UNDER-GRADUATE'S SOCIETY.

The annual dinner of the Undergraduates of the Medical Faculty of the University of Bishop's College came off on December 9, 1886, in the Ladies' ordinary of the Windsor Hotel, and was marked by much cordiality. Some seventy-five gentlemen, some representing the other universities, sat down to an excellent dinner, to which full justice was done. Dr. F. W. Campbell, M. A., M. D., L. R. C. P., London, Dean of the Faculty, occupied the On his right were seated Mr. Heneker, chair. Chancellor of the University, Dr. Anderson, United States Consul General; Dr. McEachran, Mr. J. S. Hall, M.P.P., Dr. George Ross, representing McGill, and Dr. Proudfoot. On the left of the