VII. DEXIIDÆ (Continued from page 55.)

the abdomen is rather densely set with subspinose macrochaetæ.)

- 4. Elongate subcylindrical macrotype eggs, hatching to coloured maggots in the fat gut-like uterus, maggots deposited near host-Myoceropiæ.
- 5. Very elongate subfiliform small macrotype eggs, hatching in the fat gut-like uterus to very slender subfiliform white maggots, which are deposited at weevil ovipositionpunctures in various green fruits and buds, the maggots making their way to the weevil grubs inside-Myiophasiiæ.
- Elongate subcylindrical macrotype eggs. hatching to white vellowish-white maggots cordate double-sac uterus, the maggots deposited on the food-substance-Sarcophagiæ G. U.
- 2. Elongate subcylindrical macrotype eggs, hatching to white maggots in a V-shaped double-sac uterus, the maggots deposited in the nests of various wasps and bees where they feed on stored insect food when such is present and on the early stages of the host-Metopiiæ Eumacronychiiæ, Paramacronychiiæ, Miltogrammiæ, Macronychiiæ .

VIII.. SARCOPHAGIDÆ

(1. Elongate subcylindrical macrotype eggs, hatching in a very long ir-IX. MEGAPROSOPIDÆ regular coiled uterus, the maggot clothed with long bristles-Megaprosopiæ.

N

tl

o

b he