upper third of the thigh. The femoral artery was found plugged above the line of demarcation. Signs of fatty degeneration were apparent in the anterior muscles of the thigh. Very few collateral vessels found, and those of small size. Chloroform was used for the operation; hemorrhage almost nil.

7 p.m.—Patient was given a hypodermic injection of morphia. He is doing well.

April 30th.—Patient had rather a restless night. At noon to-day he was slightly delirious. Leg was dressed antiseptically and is looking well; ordered

Bot. Bromid., - - 3iii Syrup, - - - - 3ii Aq. ad. - - - - 3vi

3ss 4, q, h.

May 1st.—Patient continued delirious till midnight; was then given a hypodermic of 3 gr. of morphia, and then slept till morning. Leg dressed antiseptically, and is doing well. Occasional intermissions in the heart's action were noticed by Dr. Bell, House Surgeon.

May 2nd.—Patient given a hypodermic of morphia last night, and slept well. Heart's action normal to-day; no intermissions noticed since.

May 4th.—Leg was dressed to-day and looks well. Patient complains of piles; suppository ordered.

R Pulv. opii, gr. $\frac{1}{2}$

Ext. Bellad. gr. 1 in each suppository.

Ordered Pulv. Glycyrrhizæ co. 3i statim.

May 5th.—Patient having been given an enema, the bowels operated this morning, not having operated since April 27th.

May 6th.—The leg was dressed to-day and looks well.

May 7th.—The patient was ordered the following ointment, as his hemorrhoids were troublesome:

B Pulv. opii, - - - gr. xx Liq. plumb. acet. - 3ii Vaseline, - - 3i

May 8th.—On opening antiseptic bandage, signs of putre-