

HEADACHE POWDERS.

Directions for use as found on the label.	Remarks.	RESULTS OF ANALYSIS.				No. of Sample.	Remarks and Opinion of the Chief Analyst.
		Weight of Contents of one powder, grains.	Acetanilide = Anti-febrin, grains.	Phenacetin, grains.	Phenazone = Antipyrene, grains.		

Concluded.

Dose for headache. One powder, repeated in thirty minutes if necessary. At least three hours should elapse before another dose is taken a seidlitz powder or other mild laxative should be taken with the first dose. Care should be taken in administering to children. Reduce dose according to age. Dose for sleeplessness one powder on retiring the powders may be taken dry on the tongue followed by a little water.	Powder No. 1	11.8	4.6	None.	None.	46017	Acetanilide declared to extent of 4½ grains. One powder contains excess. Meets requirements of P. or P. Act.
	" " 2	15.3	5.8	"	"		
Directions.—Take one powder dry on the tongue, or in a little water, and repeat in half an hour if not relieved. Take as early in the attack as possible. Do not repeat needlessly.	Powder No. 1	15.8	6.4	None.	None.	46018	Curative powers claimed. Acetanilide in excess; protected by stamp.
	" " 2	15.0	6.1	"	"		
.....	Powder No. 1	3.9	1.6	None.	None.	46019	Meets requirements of P. or P. Act. Acetanilide declared.
	" " 2	4.0	1.6	"	"		
.....	Powder No. 1	12.4	5.6	None.	None.	46020	Curative properties claimed. Meets requirements of P. or P. Act. Acetanilide declared.
	" " 2	12.1	5.4	"	"		