

Menandra raris alternis. Clayt. fl. virg. 20.

In sandy fields: New Jersey to Carolina. *¶*. July.
v. v. Flowers small.

- corymbosum*. 3. *H. exstipulatum*, erectum, ramosum, minutim pubescens; foliis alternis lanceolatis subtus cano-tomentosis, corymbis fastigiatis confertim multifloris, calycis laciniis ovatis acutis, capsulis calyce vix longioribus.—*Mich. fl. amer.* 1. p. 307.

In dry barren sand-fields, particularly along the seacoast: New Jersey to Georgia. *¶*. June—Aug.
v. v. Flowers very small, sometimes without petals; from which circumstance this and the foregoing species have been mistaken for *Lechea major*.

- rosmarinifolium*. 4. *H. exstipulatum*, stricte-erectum, simplex, canescens; foliis linearibus margine revolutis subtus cano-tomentosis, racemulis axillaribus interrupte confertifloris, calycis laciniis ovatis acutis, petalis calyce triplo longioribus.

In pine-barrens: Georgia. *Enslen.* *¶*. July. *v. s.*
in *Herb. Enslen.* Flowers bright yellow.

- carolinianum*. 5. *H. exstipulatum*, hirsutum, erectum; foliis alternis oblongo-ovalibus subdenticulatis, imis obovalibus utrinque hirsutis, pedunculis terminalibus paucis calycibusque villosissimis, calycis laciniis oblongis acutis petalis brevioribus.—*Mich. fl. amer.* 1. p. 307.

Cistus carolinianus. Walt. fl. car. 152.

Icon. *Vent. cels* 74.

In pine-barrens of Carolina and Georgia. *¶*. June—Aug. *v. v.* Flowers and leaves larger than in any other species.

426. HUDSONIA. *Gen. pl.* 822. *Willd. hort. berol.* 15.

ericoides.

1. *Hudsonia. Willd. sp. pl.* 2. p. 858.

Icon. *Willd. hort. berol.* 15. *Berg. act. holm.* 1778. t. 2.

In pine-barrens of New Jersey and Virginia. *h.* May, June. *v. v.* Flowers yellow. This delicate little shrub, not above six inches high, covers large tracts of sandy pine-woods in a similar manner as common heath.