

Distribution.—West Indies and Middle States, (Boisd.) ; South Carolina, (Abbot) ; Massachusetts, (Harris) ; Philadelphia, (Say) ; New York, (Emmons) ; Ohio, (Kirtland) ; Lake Superior, (Agassiz) ; Eastern Townships, (Gosse) ; Upper Canada, Rouge District, and Sorel.

Family III. NYMPHALIDÆ, Swainson.

Sub-Family I. SATYRIDI, Steph.

Genus 1, *Debis*, Doubleday ?

1. D. *Portlandia*, Boisd., *Oreas-marmorea Andromacha*, Hubner, (Pearly eyes).—Not common, July.

Distribution.—Western Prairies of Ohio, (Kirtland) ; Arkansas, but not in Pennsylvania, (Say) ; Massachusetts, (Harris) ; Eastern Townships, (Gosse) ; Rouge District.

Genus 2, *Hipparchia*, Fab. ; *Satyrus*, Boisd.

1. H. *Nephele* ? Kirby.—Common in grass-fields, July and August.

Sub-Family II. NYMPHALIDI, Steph.

Genus 3, *Limenitis*, Fab.

1. L. *Arthemis*, Drury, (Banded Purple).—Common, July and August.

Distribution.—Eastern Townships, (Gosse) ; Massachusetts, (Harris) ; Poland, Ohio, (Kirtland) ; Lake Superior, (Agassiz) ; North-West Territory, Lake Winnipeg, Lake of the Woods, Upper Canada and Arkansas, (Say) ; Sorel, Rouge District, and South side of the Gulf.

Genus 4, *Nymphalis*, Boisd.

1. N. *disippus*, Godt. ; *Mississippis*, Linn and Fab.—Common, August.

Distribution.—New York, (Environs) ; Massachusetts, (Harris) ; Ohio, (Kirtland) ; L'Orignal, on the Ottawa, (Bell).

Sub-Family III. VANESSIDI, Steph.

Genus 5, *Cynthia*, Fab.

1. C. *cardui*, Linn, (Painted Lady).—Abundant in some years, September and October.

Distribution.—Massachusetts, (Harris) ; New York, (Emmons) ; Rochport, Ohio, (Kirtland) ; South shores of the Gulf and New Brunswick, (R. Bell) ; Sorel, (Eu.)

2. C. *Huntera*, Fab., (Hunter's Cynthia).—Abundant on the Mountain, October 1850, (Billings).

Distribution.—Massachusetts, (Harris) ; Rochport, Ohio, (Kirtl-
l and)