

Postnotum subopaque, pollinose. Pleura reddish brown, subglabrous with a faint, grayish sheen. Halteres pale, club infusate. Legs slender, sordid yellowish brown; pilosity short, grayish; coxæ and basal part of femora yellowish, tarsi fuscous. Wings light gray, costal cells fuscous, stigma elongate, dark brown, apical portion of wing infusate, veins *Cu* and *Cu*<sub>2</sub> seamed with fuscous. Venation as in *Limnophila terra-novæ*\*\* Alex.

Abdomen yellowish brown, with a fine, pale pubescence; venter yellowish. Ovipositor brown, valves slender, curved upwards, apices pale.

*Holotype*.—♀, Eastern Harbour, Cape Breton Islands, July 17th, 1917. (A. G. Huntsman).

A very close ally of *L. terra-novæ*, with the description of which it agrees in all points, save that the cord and cross-veins are not seamed with fuscous; the *Rs* is longer than cell 1st *M*<sub>2</sub>—equal in *terra-novæ*—and subangulate at the base, with a short stump of a vein. Cell *R*<sub>2</sub> is very narrow in its basal half, whereas it widens from the base in *terra-novæ*.

#### *Tipula huntsmaniana*, n. sp.

Very similar to *Tipula angulata* Loew. Flagellar joints of antennæ bicoloured. The pale, angulate fascia incomplete, scarcely reaching the middle of cell *M*<sub>3</sub>; base of wing yellowish.

*Male*.—Length 13 mm.; wing 13.5 mm.

Head dull yellow; palpi yellowish brown, last joint dark brown and shorter than the preceding joints together. Frontal prolongation and rostrum concolorous, with a light, whitish pollinosity. Nasus short. Antennæ of moderate length, the three basal joints yellowish, joints three to ten yellowish, blackish at the base, outer joints brown. Frontal tubercle with impressed longitudinal line. Occiput with pale, yellowish-gray hairs.

Thorax yellowish gray, dull. Pronotum with dark, median spot. Präscutal stripes darker brown, ill defined; the median stripe narrowed posteriorly with a not sharply limited, median paler line, scutum dull brown, posterior border paler; scutellum and postnotum luteous, with ill-defined, median darker line. Pleura yellowish white, subsericeous; pleuro-dorsal membrane dull yellow. Halteres pale, club dark brown. Legs slender, dull, yellow; pilosity very short, blackish; coxæ yellowish white, sericeous; femora and tibiæ infusate at the tip, the latter a trifle longer than the metatarsus; tarsi fuscous. Wings with the pattern of *T. angulata*, but the pale, angulate fascia becomes narrowed in cell *M*<sub>2</sub> and does not extend beyond the middle of the latter cell.

Basal tergites of abdomen yellowish, becoming yellowish brown to fuscous posteriorly, with darker, median vitta; lateral margin of tergites paler. Venter similar to tergum. Eighth sternite rounded. Hypopygium yellowish brown; ninth tergite subquadrate with U-shaped median emargination, the lateral margin of the latter ends in a sharp point. Pleural suture entire; the pleurite transversely oval, the outer appendage narrow, erect, pointed, curved somewhat backward, hairy; the inner appendages consist of perpendicular plates, broadly rounded basally, narrowed upwardly and ending just below the free margin of the ninth tergite; ninth sternite with a deep, U-shaped emargination.

\*\*Journal New York Entomological Society, vol XXIV, Pl. 8, fig. 7.