

THE BEST ANTISEPTIC.

FOR BOTH INTERNAL AND EXTERNAL USE.

LISTERINE.

Formula.—Listerine is the essential antiseptic constituent of Thyme, Eucalyptus, Baptisia, Gaultheria and Mentha vensis, in combination. Each fluid drachm also contains two grains of refined an 1 purified Benzo-boracic Acid.

Dose.—Internally: One teaspoonful three or more times a day (as indicated), either full strength, or diluted, as necessary for varied conditions.

LISTERINE is a well known antiseptic agent—an antizymotic—especially useful in the management of catarrhal conditions of the mucous membrane, adapted to internal use, and to make and maintain surgical cleanliness—asepsis—in the treatment of all parts of the human body, whether by pray, irrigation, atomization, or simple local application, and therefore characterized by its particular adaptability to the field of

PREVENTIVE MEDICINE—INDIVIDUAL PROPHYLAXIS.

Physicians interested in LISTERINE will please send us their address, and receive by return mail our new and complete pamphlet of 36 quarto pages, embodying:

A Tabulated Exhibit of the action of LISTERINE upon inert Laboratory Compounds;
Full and Exhaustive Reports and Clinical observations from all sources, confirming the utility of LISTERINE as a General Antiseptic for both internal and external use; and particularly

Microscopic Observations, showing the comparative value and availability of various antiseptics in the treatment of Diseases of the Oral Cavity, by W. D. MILLER, A.B., Ph.D., D.D.S., Prof. of Operative and Clinical Dentistry, University of Berlin, from whose deductions LISTERINE appears to be the most acceptable prophylactic for the care and preservation of the teeth.

Diseases of the Uric Acid Diathesis.

LAMBERT'S

LITHIATED HYDRANGEA.

Kidney Alterative—Anti-Lithic.

Formula.—Each fluid drachm of "Lithiated Hydrangea" represents thirty grains of FRESH HYDRANGEA and three grains of CHEMICALLY PURE Benzo-Salicylate of Lithia. Prepared by our improved process of osmosis, it is INVARIABLY of DEFINITE and UNIFORM therapeutic strength, and hence can be depended upon in clinical practice.

Dose.—One or two teaspoonfuls four times a day (preferably between meals).

Urinary Calculus, Gout, Rheumatism, Bright's Disease, Diabetes, Cystitis, Hæmaturia, Albuminuria, and Vesical Irritations generally.

WE have had prepared for the convenience of Physicians Dietetic Notes, suggesting the articles of food to be allowed or prohibited in several of these diseases.

These Dietetic Notes have been bound in the form of small perforated slips for Physicians to distribute to their patients. Mailed gratis upon request, together with our latest compilation of case reports and clinical observations, bearing upon the treatment of this class of Diseases.

Lambert Pharmacal Company,

SAINT LOUIS, U. S.

RHEUMATISM.

Dietetic Note.—A fruit and vegetable diet is most favorable for patients with chronic rheumatic troubles.

ALLOWED.—Beef and mutton in moderation, with horse radish as a relish; fish and eggs, green vegetables and fruit, especially lemons. The skimmed milk diet has been advocated by some authors.

AVOID.—Starchy and saccharine food; all malt liquors, wines and coffee.

LISTERINE AND LITHIATED HYDRANGEA

MAY BE OBTAINED FROM

BROWN & WEBB, of Halifax, or from our Canadian Agent, W. LLOYD WOOD, of Toronto. British Agents, S. MAW, SON & THOMPSON, London.