

Charlotte Harbor, Fla. From Mrs. A. T. Slosson to whose kindness I am indebted for many favours, and to whom I give my grateful thanks.

*Diviana nymphæella*, n. sp.—Expands 21 mm. Palpi very long, much exceeding head, the second member especially being lengthened, dark fuscous, whitish in front. Head blackish fuscous, collar blackish. Thorax dull gray. Abdomen lightish gray. Forewings fuscous gray lines indistinct, the basal indicated by a deepening of the dark colour on either side, outer line near outer edge and parallel with it; discal spots distinct geminate, a marginal line of black spots. Hindwings light fuscous. Charlotte Harbor, Fla., from Mrs. A. T. Slosson.

It was my intention to erect a new genus *Palatka* for this species, more especially on account of the very long erect labial palpi, but I have concluded to place it under *Diviana*, which it nearly approaches. I mention this, as I gave the MS. name, *Palatka*, to Dr. J. B. Smith, and on this account the species stands under that generic name in his lately published "List of Lepidoptera."

*Chipeta*, n. gen.—Labial palpi long, slender, arched on second member, horizontal at end, 2nd member very long, end member short; maxillary palpi small, ocelli distinct; antennæ crenate pubescent. Tongue short, but not obsolete. Forewings long, narrow, oval, 9 veins 5 and 8 wanting, 3 and 4 separate, 10 separate. Hindwings 7 veins, 2 at angle, 3 and 4 stemmed, 5 wanting, 8 distinct. Near to *Tampa* Ragonot; the generic name from Chipeta, wife of Ouray, Chief of the Ute Indians. In Dr. Smith's List of Lepidoptera the genus is under the name *Osceola*, but that is preoccupied, so I change it to *Chipeta*.

*Chipeta perlepidella*, n. sp.—Expands 22 mm. Palpi, head and thorax dark red. Abdomen yellowish fuscous, with a reddish tinge. Forewings with costa narrowly white, the line not reaching apex, the rest of the wings bright reddish, quite even in colour. Hindwings white, narrow; pointed. Fla.

*Lipographis subosseella*, n. sp.—Expands 16 mm. Head, thorax and forewings of a dull white colour, rather evenly washed with light fuscous, giving a dirty white appearance, and this all sparingly mixed with blackish scales; line of forewings indistinct, shown rather by a darkening of edges which give a blackish blotch by basal line near inner margin and an even subterminal dark rather broad band. Two discal dots also indistinct. A marginal line of indistinct black dots. Hindwings fuscous, darkest at