

broader and more approximate on the sternal side pieces. Legs rather slender, femora not toothed, terminal joint of tarsi projecting a distance rather less than the length of the lobes of the preceding joint; claws simple.

Length 1.8 mm.; width .9 mm.

Awenre, Manitoba, Sept. 8, (Criddle).

A small species belonging to the *squamulatus* group of Dietz, in which it may best follow *persimilis*, though not agreeing closely with any species of the group. The small size and lack of true scales on the upper surface distinguishes it from all but *albopilosulus*, which differs in its erect, coarser vestiture among other characters.

LECTOTYPES OF THE SPECIES OF HYMENOPTERA
(EXCEPT APOIDEA) DESCRIBED BY ABBÉ
PROVANCHER.

BY A. B. GAHAN AND S. A. ROHWER, BUREAU OF ENTOMOLOGY,
WASHINGTON, D.C.

(Continued from page 336.)

Coccophagus brunneus. Type.—Yellow label 1384. 2nd Coll. Pub. Mus., Quebec. Badly glued.

Coccophagus compressicornis. Type.—Harrington Coll.

Coccophagus pallipes. Type.—Yellow label 1389; blue 783(s). 2nd Coll. Pub. Mus., Quebec. Head gone.

Coleocentrus mellipes. Type.—Not located.

Coleocentrus quebecensis. Type.—Not in Pub. Mus., Quebec., unless under *C. pettiti* Cress.

Coleocentrus rufus. Type.—Female, yellow label 456. 1st Coll. Pub. Mus., Quebec.

Copelus paradoxus.—See *Helorus*.

Copidosoma pallipes. Type.—Harrington Coll.

Crabro aciculatus. Type.—Female, yellow label 813. 2nd Coll. Pub. Mus., Quebec. Male, allotype, without label.

Crabro niger. Type.—Female, blue-green label 852(s), yellow label 1660. 2nd Coll. Pub. Mus., Quebec.

Crabro 4-maculatus.—See *C. 4-punctatus*. Type the same specimen.