

1901. *Dajus mysidis* RICHARDSON, Proc. U. S. Nat. Mus., XXIII, p. 579.
 1905. *Dajus mysidis* RICHARDSON, Bull. 54, U. S. Nat. Mus., p. 573.
 1909. *Dajus mysidis* G. O. SÆRS, Sec. Arc. Exped. Fram. Videnskabs-Selskabet i Kristiania, No. 18, p. 9.
 1916. *Dajus mysidis* HANSEN, H. J., Danish Ingolf Exped., Crust. Malacost. 3, pt. 5, p. 208.

Not taken by the Canadian Arctic Expedition.

Localities recorded:

North America: Five places off the east coast of Baffin land, namely, lat. $73^{\circ} 48' N.$, long. $80^{\circ} 30' W.$; lat. $72^{\circ} 33' N.$, long. $71^{\circ} 30' W.$; lat. $71^{\circ} 57' N.$, long. $73^{\circ} 56' W.$; lat. $66^{\circ} 33' N.$, long. $61^{\circ} 50' W.$; lat. $64^{\circ} 56' N.$, long. $66^{\circ} 18' W.$; also Labrador.

West Greenland: This parasite has been taken on *Mysis oculata* from the following stations: Murchison sound, about lat. $77^{\circ} 30' N.$, 25 fathoms; Duck island, lat. $73^{\circ} 55' N.$, 3-15 fathoms; Kingigtok, lat. $73^{\circ} 17' N.$, Jacobshavn, lat. $69^{\circ} 13' N.$, Claushavn, lat. $69^{\circ} 05' N.$, 5-10 and 10-15 fathoms; Egedesminde, lat. $68^{\circ} 42' N.$, 20-25 fathoms; Holsteensborg, lat. $66^{\circ} 56' N.$, and Brede fjord, about lat. $61^{\circ} 20' N.$

East Greenland: Danmark Havn, lat. $76^{\circ} 45' N.$, on *M. oculata* in from 0-5 to 10 fathoms; at Sabine island; Hurry inlet, lat. $70^{\circ} 50' N.$, Angmagssalik, lat. $65^{\circ} 30' N.$, on *M. oculata* in 10-15 fathoms.

Europe: North of the New Siberian islands; from Kostin Shar, southwest coast of Novaya Zemlia on *M. oculata*; also from Franz Josef land on the same host; Vardo and Vadso, Finmark, on the same host; Norway, lat. $66^{\circ} N.$, on *M. mixta*; on *M. oculata* at Jan Mayen.

Family Cryptoniscidae.

Clypeoniscus meinerti Giard and Bonnier.

1893. *Clypeoniscus meinerti* GIARD AND BONNIER, Bull. Scientifique de la France et de la Belgique (4), XXV, p. 421-436-444.
 1905. *Clypeoniscus meinerti* RICHARDSON, Bull. 54, U. S. Nat. Mus., p. 577.
 1916. *Clypeoniscus meinerti* HANSEN, H. J., Danish Ingolf Exped. Crust. Malacost. 3, pt. 5, p. 213.

Not taken by the Canadian Arctic Expedition.

Localities recorded:

West Greenland: Infesting *Synidotea nodulosa* Krøyer, partly from Godhavn, lat. $69^{\circ} 14' N.$, partly without special locality, but probably from a more southern locality.

Europe: Jugor Sehar, entrance to the Kara sea on *S. nodulosa*; on *Pleuroprius murchisoni* either from Iceland or the Faeroes.

SUBORDER GNATHIDEA.

Family GNATHIIDAE.

Genus *Gnathia* Leach 1814.

Syn. *Ancus* Risso (male).

Syn. *Pranzia* LATR. (larva).

Gnathia elongata (Krøyer).

1849. *Ancus elongatus* KRØYER, Voy. en Scand., Crust., pl. 30, figs. 3 a-g.
 1846-49. *Ancus elongatus* KRØYER, Naturh. Tidsskr. (2), II, pp. 388-394.
 1887. *Ancus elongatus* HANSEN, Dijnphma-Togetete Zool.-bot. Udbytte, p. 205.