

Fraser River, B.C., 1898, (*J. R. Anderson.*) At an altitude of between 5,500 and 6,000 ft. on nearly all mountains north and south of Chilliwack River, B.C. Nos. 54,434, 54,435, 54,436. Growing generally with *Parnassia fimbriata* and *Leptarrhena pyrolæfolia*. When growing the flowers are bright plum color shading to pink and creamy white. (*J. M. Macoun.*)

CASTILLEIA RUPICOLA, Fernald.

First collected in Canada, by Dr. James Fletcher and Mr. J. R. Anderson on Mt. Cheam, Lower Fraser Valley, B.C. Not rare, generally on inaccessible cliffs, both north and south of the Chilliwack River. Nos. 54,437, 54,441 and 54,438. (*J. M. Macoun.*)

CASTILLEIA ANGUSTIFOLIA, Don. var. BRADBURI, Fernald.

Confounded with *C. miniata* in our herbarium. Parson's Farm and other places near Victoria, Vancouver Island, No. 714; Nanaimo, Vancouver Island, No. 715; Deer Park, Columbia River, B.C., No. 17,277 (*Macoun.*) Thompson River, B.C. (*A. J. Hill.*) North Arm, Burrard Inlet, B.C.; on gravel bars, Chilliwack River, B.C., No. 54,439; north of Chilliwack Lake, B.C., No. 54,440. (*J. M. Macoun.*)

DASYSTOMA FLAVA, Wood.

In woods at Queenston Heights, Ont. This species seems to be confined to the Niagara region where it has been found by several collectors in recent years.

ANYCHIA CANADENSIS, (L.) B. S. P.

Amherstburgh, Ont., 1901. (*John Macoun.*) The specimens referred to *A. dichotoma* in Part x of these contributions are this species.

SALIX PRINOIDES, Pursh.

Below Brock's monument, Queenston Heights, Niagara, Ont., 1901. (*John Macoun.*) New to Canada.

ARCEUTHOBium PUSILLUM, Peck.

On *Picea nigra*, in a bog at Tracadie Beach, Prince Edward Island, July 31st, 1901. (*J. R. Churchill.*) Also collected on P. E. I. by Mr. L. W. Watson.