

line much as in *apicosa*; discal spots smaller than in its ally; posterior line straighter, without the prominent inward curve below the reniform, neither followed by a carneous shade at this place as in *apicosa*. The posterior line is followed by two fine ochrey shade lines; beyond these the subterminal space is blackish, cut by pale scales on the veins. Subterminal line fine, pale, irregular. Terminal space dusky, no apical spot; terminal margin more angulated than in *apicosa*, being produced opposite veins 3 and 4; the difference in shape of wing is quite marked. Terminal line black, interrupted, preceded by a fine edging of pale scales. Beneath much as in its ally, the discal dot on secondaries larger. *Expanse* 28 mil.; Penn., Mr. W. H. Stultz; one male specimen.

*Eustrotia secta*, n. s.

A small species with the ornamentation of primaries recalling *synochitis*. Hind wings dark fuscous with paler fringes. Fore wings shaded with whitish gray at base; median space brown below the median vein and here shaded with black, so that here a darker internal patch is formed, reminding one of the green patch in *synochitis*. Orbicular wanting; reniform two superimposed black points. Subterminal line diffuse, perpendicular, twice outwardly exerted, brownish. A dotted ferruginous terminal line; fringes fuscous gray, interlined. Three black ante-apical costal dots, preceded by pale points. Lines indistinct, anterior outwardly arcuate. *Expanse* 18 mil. *Hab.* Massachusetts, Mr. Roland Thaxter, No. 16.

*Oncocnemis aterrima*, n. s.

Fore wings dead black with obliterate ornamentation. The small reniform may be distinguished ringed with white. The inner line is lost; outer line white, evenly curved opposite the cell, slightly bent inwardly below median vein. The subterminal space is washed with white behind the outer line and this portion of the wing is the most prominently ornamented. A series of whitish points indicates the subterminal line. There is a slight olivaceous powdering over the primaries and thorax. Hind wings dead black, with a median curved dark line brought into relief by a following pale shading; fringes whitish. Beneath black with a common median line followed by white streaks and shading. Eyes naked; fore tibiae with a stout claw. *Expanse* 22 mil. *Habitat*, Havilah, Cal., Mr. Hy. Edwards, No. 119. This species has a slight Heliothid appearance and differs much in markings from the others known to me in this genus.