

"Secondaries without spots, sometimes with a dark spot near the external edge, and faintly marked near the external angle.

"Abdomen white, with three rows of minute black spots, frequently inconspicuous. The fore coxae and femora luteous; tarsi blackish.

"Length of body 5-8 lines. Wings expand 13-18 lines."

Larva undescribed.

Hab.—Mass.; Penn.; Georgia.

**H?* (*Spilosoma*) *punctata*.—Fitch. Fitch's Third Report, p. 265.

"Primaries white, with a black central dot, and in the males a row of small blackish spots, extending from the middle of the inner margin to the tip.

"Secondaries white.

"Thighs and hips yellow in front; a continuous black stripe on fore side of anterior feet and shanks."

Larva undescribed.

Hab.—New York.

Euchates.—Harris.

Wings bluish-gray, without spots. Abdomen smooth, spotted. Hind tibiae with two pairs of spurs.

**E. Egle*.—Drury. Figured in Drury II., pl. 20. Larva figured in Harris' Insects, new edition, fig. 172.

"Head gray; occiput with a narrow luteous line. Thorax gray.

"Wings rather long, thin, and delicate, of a bluish-gray color, paler on the front edge, and without spots.

"Abdomen above dark yellow, with a dorsal and lateral row of black spots; beneath whitish or gray. Fore coxae woolly, and touched at the sides with luteous."

Wings expand 17-20 lines.

"Larva black, with a whitish line on each side, and thickly covered with short tufts of hairs, proceeding from little warts. Along the top of the back is a row of short black tufts, and on each side, from the fifth to the tenth ring inclusive, are alternate tufts of orange and yellow hairs, curving upwards so as nearly to conceal the black tufts between them; below these, along the sides of the body, is a row of horizontal black tufts. On the first and second rings are four long pencil-like black tufts, extending over the head; on each side of the