

has run through the woods. Specimens of this beetle are often found in recently built houses and about lumber-yards.]

[169.] 223. *MONOCHAMUS MARMORATOR Kirby*.—Length of body 11 lines. A single specimen taken in Lat. 54°.

Body black, covered underneath, but so that the black appears in various places, with subcinereous, or somewhat tawny decumbent hairs. Head and prothorax covered in the same way but with redder hairs: spines of the prothorax very robust, rather long, sharpish: scutellum covered with a coat of cinereous hairs, divided by a black longitudinal line: elytra black, marbled variously with cinereous and reddish tawny hairs; the cinereous spots are dotted with black; the surface of the elytra when laid bare appears punctured, and at the base are several confluent smooth elevated spaces; suture and lateral margin testaceous; apex acute.

N. B.—The antennæ in the specimen are broken off. [Unknown to us.]

224. *ACANTHOCINUS (GRAPHISURUS) PUSILLUS Kirby*.—Length of body $4\frac{1}{4}$ lines. A single specimen taken in the journey from New York to Cumberland-house.

[170.] This species is one of the most minute of the Capricorn tribes. Body linear, black but covered with a coat of whitish decumbent hairs, which appears more or less sprinkled with black dots. Head longitudinally channelled; antennæ mutilated in the specimen, but those joints that remain are white at the base: prothorax short, armed on each side, towards the base with a short sharp spine, punctured with scattered punctures; elytra punctured especially towards the base, mottled and speckled with brown, with an oblique brown band a little beyond the middle, apex of the elytra rounded: podex and hypopygium, or last dorsal and ventral segments of the abdomen elongated, so as to defend the base of the ovipositor which is exserted, causing the insect to appear as if it had a tail; the hypopygium is emarginate: thighs much incrassated at the apex. [Not common; taken at Grimsby by Mr. Pettit, and on oak-trees in the neighbourhood of Philadelphia by Mr. Bland.]

225. *CALLIDIUM AGRESTE Kirby*.—Length of body 11 lines. Several specimens taken in the Expedition, and likewise in Nova Scotia by Dr. Mac Culloch and Capt. Hall.

I at first took this for a variety of *C. rusticum*, but on a closer inspection I found it differed in the sculpture as well as colour; and having