

WGI POLICYMAKERS SUMMARY

these organisms release carbon dioxide via respiration into the atmosphere. Most land plants have a system of photosynthesis which will respond positively to increased atmospheric carbon dioxide ("the carbon dioxide fertilization effect") but the response varies with species. The effect may decrease with time when restricted by other ecological limitations, for example, nutrient availability. It should be emphasized that the carbon content of the terrestrial biosphere will increase only if the forest ecosystems in a state of maturity will be able to store more carbon in a warmer climate and at higher concentrations of carbon dioxide. We do not yet know if this is the case.

The response to increased carbon dioxide results in greater efficiencies of water, light and nitrogen use. These increased efficiencies may be particularly important during drought and in arid/semi-arid and infertile areas.

Because species respond differently to climatic change, some will increase in abundance and/or

range while others will decrease. Ecosystems will therefore change in structure and composition. Some species may be displaced to higher latitudes and altitudes, and may be more prone to local, and possibly even global, extinction; other species may thrive.

As stated above, ecosystem structure and species distribution are particularly sensitive to the rate of change of climate. We can deduce something about how quickly global temperature has changed in the past from paleoclimatological records. As an example, at the end of the last glaciation, within about a century, temperature increased by up to 5°C in the North Atlantic region, mainly in Western Europe. Although during the increase from the glacial to the current interglacial temperature simple tundra ecosystems responded positively, a similar rapid temperature increase applied to more developed ecosystems could result in their instability.

Deforestation and Reforestation

Man has been deforesting the Earth for millennia. Until the early part of the century, this was mainly in temperate regions, more recently it has been concentrated in the tropics. Deforestation has several potential impacts on climate: through the carbon and nitrogen cycles (where it can lead to changes in atmospheric carbon dioxide concentrations), through the change in reflectivity of terrain when forests are cleared, through its effect on the hydrological cycle (precipitation, evaporation and runoff) and surface roughness and thus atmospheric circulation which can produce remote effects on climate.

It is estimated that each year about 2 Gt of carbon (GtC) is released to the atmosphere due to tropical deforestation. The rate of forest clearing is difficult to estimate; probably until the mid-20th century, temperate deforestation and the loss of organic matter from soils was a more important contributor to atmospheric carbon dioxide than was the burning of fossil fuels. Since then, fossil fuels have become dominant; one estimate is that around 1980, 1.6 GtC was being released annually from the clearing of tropical forests, compared with about 5 GtC from the burning of fossil fuels. If all the tropical forests were removed, the input is variously estimated at from 150 to 240 GtC; this would increase atmospheric carbon dioxide by 35 to 60 ppmv.

To analyse the effect of reforestation we assume that 10 million hectares of forests are planted each year for a period of 40 years, ie 4 million km² would then have been planted by 2030, at which time 1GtC would be absorbed annually until these forests reach maturity. This would happen in 40-100 years for most forests. The above scenario implies an accumulated uptake of about 20GtC by the year 2030 and up to 80GtC after 100 years. This accumulation of carbon in forests is equivalent to some 5-10% of the emission due to fossil fuel burning in the Business-as-Usual scenario.

Deforestation can also alter climate directly by increasing reflectivity and decreasing evapotranspiration. Experiments with climate models predict that replacing all the forests of the Amazon Basin by grassland would reduce the rainfall over the basin by about 20%, and increase mean temperature by several degrees.