

Fam. EUSCHEMIDÆ.

PITTHEA, Walk.

PITTHEA CONTINUA.

Pitthea continua, Walk. Cat. ii. p. 463.

One specimen, that in no respect differs from examples from Lagos, Addah, Cameroons, and the Congo in my collection. This species has a wide range in Western Africa; it has not, so far as I know, yet been recorded from East Africa, its place being taken on the East Coast by *Pitthea trifasciata*, Dewitz, Nova Acta Acad. Nat. Cur. xlvii. p. 82, t. iii. f. 3.

PITTHEA AGÉNORIA, sp. n.

Primaries and secondaries deep black, both the wings crossed by a semihyaline white band, that on the secondaries not reaching the outer margin; the underside the same as above; the head, antennæ, thorax, abdomen, and legs deep black; the collar and the underside of the abdomen bright orange-red. Expanse $1\frac{3}{4}$ inch.

Two specimens of this very distinct species are in the collection: it is allied to *Pitthea trifasciata*, Dewitz, from which it is at once distinguished by the entire absence of the apical hyaline band on the primaries, and by the anal segments being black instead of yellow as in that species.

Fam. SPILOMELIDÆ.

ZEBRONIA, Hübn.

ZEBRONIA PHENICEALIS.

Phalæna-Pyralis Phenice, Cram. Pap. Exot. iv. p. 185, t. 382. f. G.

Zebronia fæderalis, Hübn. Verz. bek. Schmett. p. 361.

Spilomela picealis, Guén. Sp. gén. des Lép. viii. p. 281.

Spilomela podatirialis, Guén. Sp. gén. des Lép. viii. p. 281.

Zebronia phenicealis, Walk. Cat. xvii. p. 476; Druce, Moloney's Forestry of Western Africa, appendix v. p. 505.

One specimen, identical with those before me from the Gambia and the Cameroons.