216. Oberea schaumii LeConte.

Found in various parts of California, more common in the Sierras; breeds in willow and cottonwood, taken by Van Dyke in the Sierras, breeding in live twigs of the latter.

217. Tetraopes elegans Horn.

Taken at San José del Cabo, Lower California.

218. Tetraopes femoratus LeConte.

Taken in Sierras. May 11-June 17. Very common at times on milkweed.

var. mancus LeConte.

Common in Southern California on milkweed.

var. basalis LeConte.

Common throughout the Sierras on milkweed.

var. oregonensis LeConte.

Found in northern Sierras on milkweed.

219. Lianema tenuicornis Fall.

Found at El Taste, Lower California.

220. Idæmea fulleri Horn.

Reported from the southern part of the State. *I.californica* is, however, the one most probably seen in this State.

221. Idamea californica Fall.

Beaten by Fall from live oak at Pasadena in June; others taken by Fall at light; several taken by Van Dyke on the sea shore at Santa Monica.

222. Styloxus lucanus LeConte.

Found in Lower California.

223. Methia æstiva Fall.

Found around electric lights at Pasadena.

224. Dysphaga debilis Horn.

Found at San José del Cabo, Lower California.

Additions to List.

225. Mallodon molarium Bates.

Found at San José del Cabo, Lower California.

226. Acyphoderes delicatus Horn.

Found at El Taste, Lower California.

There are four species in the list which were given separate numbers which may be either synonyms or varieties (84–85, 131, 156, 199). Therefore, there are 178 unquestioned species confined to California, 10 present both in California and Lower California, and 43 confined to Lower California.