

stems with the flax-seed (pupa) in the centre of the stem. The larvæ live gregariously, according to Mr. A. Costa, but I have never found more than four pupæ at one joint.

After all, it would be of prominent importance to compare the species from Naples with specimens of the true Hessian fly, which was collected there long ago by Prof. Dana.

DETERMINATION OF THE 36 COLEOPTERA DESCRIBED BY D. ZIEGLER.

BY SAMUEL HENSHAW, BOSTON, MASS.

"Descriptions of New North American Coleoptera," Proc. Acad., 1844, v. 2, p. 43-47; 1845, v. 2, p. 266-272.

	Oxyporus pulcher.	43	W. Pa.	= O. femoralis, Grav.
	Diacanthus splendens.	44	"	Belongs to Corymbites.
	Scyrtes suturalis.		Pa.	= S. orbiculatus, Fabr.
	Hydnocera? longicollis.		"	
5	Spercheus tessellatus.		"	= Belongs to Hydrobius.
	Hydrophilus ovalis.	45	"	Name = H. ovatus, G. & R.
	Coprobis obtusidens.		"	= Canthon laevis, Drury.
	Pandarus? brunneus.		"	Belongs to Anaedus.
	Cistela marginata.	46	"	
10	erythroptera.		"	= C. brevis, Say.
	Pedilus nigricans.		"	= Corphyra lugubris, Say.
	haemorrhoidalis.		"	{ = " elegans, Hentz.
	ruficollis.	47	"	
	marginicollis.		"	= " labiata, Say.
15	Monohammus tomentosus.		"	= Goes tigrinus, DeGeer.
	Oedionychis? hispida.		"	= Hypolampis pilosa, Illig.
	Philonthus ater.	266	Pa.	= Quedius capucinus, Grav.
	Tachinus puncticollis.	267	"	= T. fimbriatus, Grav.?
	Phoenops luteosignata.		Geo.	= Melanophila notata, L. & G.
20	Agrilus quadriimpressus.		"	= A. acutipennis, Mannerh.
	Limonijs definitus.	268	Car.	