

has less black, but there are two rows of black spots on all but the last segment, and is slightly darker on the lateral margins. Tibiæ darker.

From North Carolina.

*THERIOPECTES TETRICUS*.—Length 17 mm.

Female.—Eyes pubescent, ocelligerous tubercle present. Front black; callosity black, shining, with an unconnected black spot above; sub-callous black, denuded. Face and cheeks black with white hairs. Antennae red, third joint black, upper angle projecting but little. Palpi white with white hairs. Thorax grayish-black with the usual gray lines and gray pubescence; humerus reddish-brown; pleurae and pectus gray with long white hairs. Abdomen black, with three rows of white triangles, the middle row indistinct; the sides of segments two and three reddish; hind margins with a fringe of white hairs, which is wanting between the middle and outer triangles. Venter yellowish-red with gray pollen, darker towards the tip. Femora black, yellowish-brown at the tips; tibiae brown, darker on the distal half; tarsi black. Wings hyaline.

From Montana.

*THERIOPECTES FRENCHII*.—Length 14 mm.

Female.—Front black; callous dark brown with a line extending above; sub-callous black. Face and cheeks black with white hairs; palpi yellowish with minute black hairs. Antennae red, annulate portion of third joint black, angle not prominent. Thorax black with the usual lines; humerus reddish-brown; pleurae and pectus with white hairs. Abdomen black with three rows of triangles, the lateral rows from segments one to four prominent, middle row largest on second segment, on the other segments only an expansion of the hind margins; lateral margins of segments 4-7 yellow with white hairs. Venter reddish-yellow, darker towards the tip, covered with gray pollen. Femurs black, with whitish hairs, brown at the tips; tibiae brown, with white hairs, darker at the tips; tarsi black. Wings hyaline with faint clouds on the cross veins and bifurcation of third vein.

From Montana.

*THERIOPECTES SUSURRUS*.—Length 14 mm.

Female.—Front gray; callosity brownish-black with black line above; ocelligerous tubercle brown. Face and cheeks white with white hairs; palpi yellow with minute white hairs, and a few black hairs. Antennae