

half their length beyond the apex of the plates, ligulate in form and curved downward and outward at apex; pygofers rounded at apex, almost equalling the plates, the small anal aperture placed superiorly.

New York; Connecticut. Described from three males. One taken at Portage Falls, N. Y., May 30th, 1888. The other two specimens were taken by me in Connecticut in the spring of 1883. One of these was swept from weeds and bushes on the hills about Northford, June 26th; the other, a very deeply coloured example, I found in a grove on Prospect St. in New Haven, June 4th. The long extruded styles of the males will at once distinguish this from any other of our described species of *Athysanus*.

5. *ATHYSANUS INSTABILIS*, n. sp.

More elongated than most of our species of this genus. Black, irrorated and finely marked with pale yellow. Elytra pale, the areoles edged with fuscous. Length, 4-4½ mm.

Head obtusely pointed before; vertex about ⅓ longer on the middle than next the eye, passage to the front well rounded, base with a few impunctured areas. Ocelli large, fulvous, distant from the eyes; temples broad; front convex, the sutures arquated at the antennæ and incurved at apex; clypeus feebly narrowed towards its slightly rounded apex; cheeks rather narrow, but feebly angled, not exceeding the clypeus. Pronotum rather long, with the sides long and carinated, latero-posterior edges straight, angles prominent, obtuse, surface strongly punctured or shagreened. Elytra long, with the appendix well developed. Valve of the male short, obtuse at apex; plates forming an equilateral triangle, fulvous clouded with dusky, their edges nearly straight, fringed with pale bristles. Last ventral segment of the female slightly produced and rounded on the middle, feebly arquated either side within the short subacute lateral angles; pygofers rather small, slightly exceeded by the oviduct.

Colour black. Base of the vertex, outer edge of the cheeks, and margins of the eyes slenderly yellow. Two transverse spots on the disc of the vertex near its base, two lines anterior to these, one or two dots near each eye, an angled slender line on the apex including a dot behind, a dot on the temples, about six arcs, a basal spot and a longitudinal line on the front, the latter expanded against the base of the clypeus, and