## HEADACHE POWDERS

		1					
Directions for use as found on the label.	Remarks.	RESULTS OF ANALYSIS.					
		Weight of Con- tents of one pow- der, grains.	Acetanilide= Anti-febrin, grains.	Phenacetin, grains.	Phenazone=Anti- pyrine, grains.	No. of Sample.	Remarks and Opinion of the Chief Analyst.
R. J. WAUGH, INSPI	ECTOR.						
Soak the wafer in water until soft, then place in the mouth and swallow with drink of water. The dose may be re- peated in half an hour if necessary. For rheu- matic pains. Take one wafer three times a day.	n n 2		None.	None.	None	.42276	Apparently contains aspirin (aceto-salicy-lic acid). Meets requirements of P. or P. Act.
Directions.— Children under 10 years give \( \frac{1}{2} \) of a powder; ten to fifteen years, \( \frac{1}{2} \) a powder, repeated in half an hour if needed. Place powder on the tongue and take a swallow of water.	2		3.3		H	42277	Acetanilide declared Meets requirements of P. or P. Act.
Directions,—Take one powder dry on the ton- gue or in a little water, and repeat in half an hour if not relieved. Take as early in the attack as possible, do not repeat needlessly.	и и 2	. 8 5 8 5	1.7			42278	Claim curative effects otherwise meets re- quirements of P, or P, Act.
	Powder No. 1	4·7 5·6	None	None.		42279	Apparently contains as pirin (aceto-salicyli acid) needs require
	Powder No. 1	5.1			None.	42280	ments of P. or P. Act Meets of requirement of P. or P. Act