

DR. A. C. DANIELS' MEDICINES

can obtain Daniels' Colic Cure you may give *two doses of each* (60 drops) instead of one to start with, then proceed as above, or giving once in five to ten minutes apart. Usually four to eight doses will cure a bad case, and two to four a mild case. Keep everything as quiet as possible. If the horse finds any position in which he seems easier, let him keep it. If convenient give him an injection of blood-warm soapsuds, or introduce the hand, well greased, into his rectum and remove all excrement possible. Many times he will be found so constipated that the gas cannot work off naturally. Keep cool. Don't get nervous or give other medicines. This remedy will do the work, and never fails if given in any reasonable season. It has cured thousands of cases after they had been given up as lost by good veterinarians. "While there's life there's hope," even if the eyes are filmed, his upper lip turned up, his pulse cannot be felt, and you can prick him with a pin without his moving. Don't despair. He may recover even when as near death as that. When the horse is cured of colic, give a few doses of Dr. Daniels' Fever Drops, two hours apart, to guard against inflammation of the bowels which is almost sure to follow a long-continued case of colic.

Paralysis of Hind Quarters

This trouble has become very common among fine horses, and is greatly dreaded by horsemen. It usually attacks horses that are in fine condition, that have not had their customary amount of exercise for a few days. The horse comes from the stable in fine spirits, but in an hour or two grows stiff in one or both hind legs; soon falls flat, and is unable to rise, or sits upon his haunches like a dog, and is perfectly helpless. The No. 1 Colic Drops have been used in many cases with good success; and if you have no good veterinarian, proceed as follows:

Treatment

Give thirty drops of No. 1 every fifteen minutes for two or three hours; later once in thirty minutes. After he shows signs of improvement, gradually reduce the dose. Wring woolen blankets out of as hot water as can be borne, and folding them, cover the back and hips, changing them often. Cover with a rubber blanket to keep in the steam. Rub hind legs with strong mustard water. Sling him up if you can; if not, keep deep, soft bedding under him, and turn him over and change his position twice in twenty-four hours anyhow; four times is better.

Feed warm bran mashes. Mix into it two of Dr. Daniels' Renovator Powders each day. Avoid all noise. Keep him as easy and comfortable as you can in every way. If attended to in first stages, this treatment is generally successful.

Stoppage of Water and Blackwater

Treat as for colic in first stages, also give a few doses of my Renovator. If these do not relieve in two hours, the water must be drawn with a catheter.

Colic, Hoven or Bloat in Cattle

Give full-grown cattle 30 drops Colic Drops on the tongue every ten minutes; 3 to 5 doses are usually required. Calves and yearlings, 5 to 10 drops is a dose. Feed my Renovator occasionally and prevent this trouble.

Scours. — Give No. 1 Colic Drops once in half hour. A few doses will generally check it.