

I. not quite twice as long as femur III., metatarsus I. barely longer than tibia I.; abdomen higher than long, globose; region of epigynum swollen; there is a small median triangular black projection or finger.

Brazos Co., Texas; Sept.

*Plasiocrærius lobiceps*, n. sp.

Length 1.5 mm. Cephalothorax yellowish with a black margin, each eye with a black ring, a black line on each side of the lobe; mandibles yellowish, legs and palpi yellowish, sternum red, black on margins; abdomen black, spinnerets pale. Head of male moderately elevated into a large lobe, bearing the P. M. E., which are large and scarcely twice their diameter apart; a hole on each side just behind the S. E.; the mandibles show a series of transverse lines on the outer side; legs moderately long, first pair longest, no spines above on the tibiae; sternum broad, triangular, bluntly pointed behind. Male palpi quite long; the tibia with a broad extension above and a hook on the inner side; the tarsus short, truncate at tip; the bulb, in side view, is constricted near the middle, the upper part crossed by two transverse dark lines, the black style coiled around the tip once, a small triangular hook near base of bulb. In the female the head is scarcely elevated; the epigynum shows a semicircular area limited by a concave ridge in front, from which there extends behind a gradually broadening furrow with its margins at tip, curved outward and backward.

One from Chicago, Ill., under leaves in October; others from Salineville, Ohio. (A. D. MacGillivray.)

*Icius canadensis*, n. sp.

Length ♀ 5 mm.; ceph. 2.4 mm. long, 1.9 mm. broad; tibia plus patella IV. 2 mm. The male but little smaller. Cephalothorax red-brown, black in eye-region; mandibles reddish; leg I. reddish except the yellowish tarsi, other legs wholly pale yellowish. Sternum infuscated; abdomen brownish with a narrow white line around base, and pale chevrons toward tip, venter pale gray, with a straight jet black stripe each side, and a narrow basal median spear-mark; a black spot each side at base of spinnerets; in ♂ more white hair around the A. M. E. Eye-region one and a fourth broader than long, broader behind than in front, first eye-row curved; eyes of second row half way between dorsal and lateral eyes; cephalothorax moderately high; mandibles vertical, with one stout tooth on inner edge of fang-groove. Legs moderately long, IV. pair longest, I. pair very stout, three pairs of spines on the tibia and