

*SELIDOSEMA LACHRYMOSA*, n. sp.

Expands 30 mm. Palpi black, tipped with ochreous; front black; thorax blackish fuscous; abdomen blackish fuscous. All wings nearly uniform blackish fuscous, fore wings rather narrow, apex acute, outer margin rounded, inner margin long; inner line well out, rounded, a sinus at cell opposite discal spot, and a less one at vein 2; outer line beginning on costa near apex, strongly sinuous, the largest sinus outwardly at vein 3, and a short, almost angular one close to inner margin, whose middle the line reaches, or a little beyond it; the lines are fine, black, the outer faintly edged outwardly with gray; a faint whitish, sub-wavy, submarginal line nearly parallel with outer margin; discal spot rather large, oval, white. Hind wings triangular, both angles prominent and sharp, outer margin irregularly wavy; a faint discal shadow line; an outer fine, black, somewhat undulating line; discal spot distinct, white; marginal lines on all wings fine, black, broken. Beneath almost even blackish fuscous, smoother than above.

Los Angeles Co., Cal.; taken in July.

*CLEORA SUBAUSTRALIS*, n. sp.

Expands 42 mm. Palpi moderate, ascending, fuscous brown, black in front and at end; front dark smoky fuscous; summit fuscous brown; thorax dark fuscous, with a bluish tinge, behind lighter; abdomen ochre fuscous, stained and dotted dorsally with blackish, incompletely interlining the segments. Wings light brownish ochre, heavily and quite evenly overlaid with blackish patches and striations, giving a generally mottled appearance, enough separated basally and outwardly to give faint indications of rounded sinuate lines of ground colour; a lighter spot outwardly at vein 3; a large, lengthened, black discal spot, and a marginal line of rather heavy intervenular black points. Hind wings corresponding to fore wings, a lighter spot at vein 3, a faint zigzag lighter outer line, a lengthened black discal spot, and a line of black intervenular lunules. Beneath even light ochre brown; fore wings darkened basally along costa, and outwardly below apex; discal spots on all wings, long, black; some faint blackish striæ scattered on fore wings.

Cocoanut Grove, Florida. National Museum collection. The generic reference is provisional. Type No. 3960.

*CLEORA PEDICELLATA*, n. sp.

Expands 43 mm. Palpi rather long, porrect, rather slender, ochre, heavily marked with black, last segment black; front tufted, ochre