ı.	Abdomen and thorax with the hair entirely black
	(Texas) melanosoma, Ckll.
	Abdomen with conspicuous white hair at sides, at least
2.	Hair of pleura and mesothorax largely dull white (S. Illinois,
	Robertson) bimaculata, Lep., var. a.
	Hair of pleura and mesothorax black 3.
3.	Mandibles with a large yellow spot (Baldwin, Kansas, July,
	Bridwell) bimaculata, Lep., var. b.

Mandibles with at most a very minute yellow dot (Ames, Iowa,

Anthedon compta (Cresson).

Both sexes of this magnificent species were taken at Fedor, June 19, 1899. It is new to the fauna of Texas.

Anthophora abrupta, Say.

Fedor; the female, April 8, 1904; males, April 27 and 29. Unless the venation is examined, this will be likely to be confused with Emphoropsis floridana Fedorensis.

BOOK NOTICES.

A CATALOGUE OF THE ERYCINIDE OF THE WORLD.—By Levi W. Mengel, Professor of Natural History, Boys' High School, Reading, Pa. I vol., pp. 161. (Price \$2.00.)

This very full and comprehensive work will be of great value to all students of Butterflies who do not confine their attention to the species inhabiting their own country. It is similar in arrangement and style to Dr. Skinner's well-known Catalogue of North American Rhopalocera, giving full bibliographical references and habitat for each species. Its extent may be realized by the following comparison: In the genus Libythea Dr. Skinner gives 2 species and Prof. Mengel 21; in the sub-family Lemoniinæ the former has two genera, including 11 species, the latter 86 genera and an enormous number of species. The book is very clearly printed and is made complete by a full index of all the species and synonyms contained in it. It may be obtained from the author.