

17th, 1887. Collected July 17th, 1887, on Mount Arrow-smith, V.I., and May 19th, 1893, on Parson's Mountain near Victoria, V.I., by Prof. Macoun. The specimens collected in 1887 formed part of the material upon which Watson based his *S. occidentalis*.

*SAXIFRAGA LATA*, (Small.)

*S. occidentalis*, Cat. Can. Pl., II: 321 in part.

*Micranthes lata*, Small, N. A. Fl., XXII: 145.

Described from specimens collected by Prof. John Macoun at Lytton, B.C., April 16th, 1890. Not rare west of the coast range. ‡

*RIBES OXYCANTHOIDES*, L. var. *CALCIOLA*, Fernald, *Rhodora*, VIII: 155.

This variety resembles the species "but young branches, petioles and lower leaf surfaces permanently and densely white-tomentose." Collected or noted growing in calcareous soils in Bonaventure and Gaspé counties, Quebec, by Messrs. Collins, Fernald and Pease at the following places: Carlisle, Tracadigash Mt., mouth of Bonaventure River, New Richmond, Grand River, Percé, Little Cascapedia River and Dartmouth River. This variety was collected on "the island" Baddeck, Cape Breton Island, N.S., by Prof. Macoun in 1898, Herb. No. 19,102. An intermediate form was collected on Cape Breton Island, the same year at Grand Narrows, No. 19,100, and Big Intervale, Margaree, No. 19,101.

*CRATAEGUS BRUNETIANA*, Sargent.

A common species about Quebec. Our specimens were collected in fruit near Quebec in 1902 by Dr. Robt. Bell.

*SPIRÆA SORRIFOLIA*, L.

Escaped from cultivation and well naturalized on the bank of the Gatineau River at the railway station, Wakefield, Que., 1903. (*John Macoun*)

*GEUM PULCHRUM*, Fernald, *Rhodora*, VIII: 11.

Known in Canada only from specimens collected by Williams, Collins and Fernald, in boggy meadows by the St.

‡The Geographical limits given in these papers refer to Canada only.