

sexes, darker outwardly; in the male the wing is somewhat lighter towards base than in ♀; discal dot and mesial band apparent, though not prominent. Fringe white. Beneath primaries light yellowish-brown. Blackish blotch on middle of costa and two at outer end, the last two marking the beginning of obscure dusky bands across the wing. Discal dot indicated, though not prominent, fringes as above. Secondaries dusky along costa and external margin, otherwise whitish. Quite well marked median band and discal dot. Body parts beneath, whitish. Legs whitish, checkered and shaded with black.

Type: 2 ♂, 1 ♀, So. Arizona, Pinal and Pima Co. Mr. Poling.

*Antaplagia hachita*, n. sp.—Expanse: 23 mm.

The space between t. a. and t. p. lines creamy white, remainder of fore wings bright chrome yellow. Transverse lines black, distinct. T. a. almost transverse, with slight outward projections below costa, above inner margin and on median vein. T. p. slightly incurved below costa, moderately exerted beyond cell, thence only slightly oblique to inner margin. The t. a. and t. p. lines are in a general way parallel, and divide the wing quite accurately into thirds. A row of black spots runs through the middle of the terminal third, following line of outer margin, the second and third from costa subobsolete, the others gradually decreasing in size towards inner margin. Fringe pale yellowish. Hind wings slightly yellowish fuscous, darker outwardly, very faint trace of mesial band, especially at costa. Beneath, fore wings narrowly yellow along costa, more broadly along outer margin, the remainder blackish, with evident mesial band corresponding to t. p. line above. Hind wings pale yellowish-white. Head and collar chrome yellow, thorax rubbed, abdomen greased. Front of head with crater-like protuberance.

Type: 1 ♀, Santa Catalina Mts., Ariz., August. Mr. Poling.

*Stibadium ochoa*, n. sp.—Expanse: 30 mm.

Ground colour pale yellow, tinged with green of a light olivaceous tint. Lines and spots marked with a darker olivaceous green or castaneous shade, which shade also covers the lower part of the median space, as well as between the ordinary spots. The basal part of the wing, as well as along costa, is quite thickly sprinkled with violet scales, and less profusely over median and subterminal spaces. Palpi dark castaneous. Head dark iron-gray. Collar and thorax castaneous, concolorous with fore wing. Abdomen in both specimens greasy, but, as near as can be told, of a somewhat more yellowish tinge than thorax. Fore wing with