TABLE III.
For five years ending 31st March, 1838.
Aggregate strength of the troops................15,674
Average annual strength......................... 3,135
Number of cases............................. 59
Number of deaths............................. 5

Ratio of cases to strength as 1 to 265.

Ratio of deaths to cases as 1 to 12.

TABLE IV.
Five years ending 31st March, 1843.
Aggregate strength of the troops................56,188
Average annual strength.......................11,237
Number of cases............................. 411
Number of deaths............................. 36

Ratio of cases to strength as 1 to 136.

Ratio of deaths to cases as 1 to 11.

TABLE V.
Five years ending 31st March, 1848.
Aggregate strength of the troops................35,567
Average annual strength......................... 7,113
Number of cases............................. 59 .
Number of deaths............................. 41

Ratio of cases to strength as 1 to 60.

Ratio of deaths to cases as 1 to 14.

TABLE VI.
Five years ending 31st March, 1853.
Aggregate strength of the troops................27,912
Average annual strength......................... 5,582
Number of cases............................. 509
Number of deaths............................. 50

Ratio of cases to strength as 1 to 54.

Ratio of deaths to cases as 1 to 10.

RECAPITULATION.
Number of cases of Delirium tremens in thirty years............ 1769
Number of deaths....................................... 143

Ratio of cases to strength, first fifteen years, as 1 to 175.

Do.,   second fifteen years, as 1 to 75.

Do.,   the whole thirty years, as 1 to 93.

Ratio of deaths to cases, first fifteen years, as 1 to 16.*

Do.,   second fifteen years, as 1 to 12.

Do.,   the whole thirty years, as 1 to 12½.

* Vide Note to Table I.